

An exploratory study of the neural mechanisms of decision making in compulsive hoarding

D. F. Tolin^{1,2*}, K. A. Kiehl³, P. Worhunsky¹, G. A. Book¹ and N. Maltby¹

¹ The Institute of Living/Hartford Hospital, Hartford, CT, USA

² Yale University School of Medicine, New Haven, CT, USA

³ MIND Institute and University of New Mexico, NM, USA

Background. Prior studies have suggested unique patterns of neural activity associated with compulsive hoarding. However, to date no studies have examined the process of making actual decisions about whether to keep or discard possessions in patients with hoarding symptoms. An increasing body of clinical data and experimental psychopathology research suggests that hoarding is associated with impaired decision making; therefore, it is important to understand the neural underpinnings of decision-making abnormalities in hoarding patients.

Method. Twelve adult patients diagnosed with compulsive hoarding, 17% of whom also met criteria for obsessive–compulsive disorder (OCD), and 12 matched healthy controls underwent functional magnetic resonance imaging (fMRI) while making decisions about whether or not to discard personal paper items (e.g. junk mail) brought to the laboratory as well as control items that did not belong to them. Items were either saved or destroyed following each decision.

Results. When deciding about whether to keep or discard personal possessions, compulsive hoarding participants displayed excessive hemodynamic activity in lateral orbitofrontal cortex and parahippocampal gyrus. Among hoarding participants, decisions to keep personal possessions were associated with greater activity in superior temporal gyrus, middle temporal gyrus, medial frontal gyrus, anterior cingulate cortex, precentral gyrus, and cerebellum than were decisions to discard personal possessions.

Conclusions. These results provide partial support for an emerging model of compulsive hoarding based on complications of the decision-making process. They also suggest that compulsive hoarding may be characterized by focal deficits in the processing of reward and changes in reward contingencies, particularly when these are perceived to be punishing.

Received 2 August 2007; Revised 3 March 2008; Accepted 15 March 2008; First published online 19 May 2008

Key words: Decision-making, fMRI, hoarding, obsessive–compulsive disorder.

Introduction

Compulsive hoarding is characterized by (a) acquisition and failure to discard a large number of possessions; (b) clutter that precludes activities for which living spaces were designed; and (c) significant distress or impairment in functioning caused by the hoarding (Frost & Hartl, 1996). Hoarding has been associated with impairment in activities of daily living (Frost *et al.* 2000), substantial health risks (Steketee *et al.* 2001), and marked occupational and role impairment (Tolin *et al.* 2007).

Although commonly considered a subtype or dimension of obsessive–compulsive disorder (OCD), a

large percentage of individuals who hoard display no other OCD symptoms (Frost *et al.* 2006), and hoarding may be as prevalent in patients with other anxiety disorders as it is in patients with OCD (Meunier *et al.* 2006). Factor and cluster analyses indicate that hoarding consistently emerges as a distinct symptom type (see Calamari *et al.* 2004, for a review). Although various OCD symptoms seem to be closely related to one another, hoarding does not seem to be particularly closely associated with OCD and is just as closely related to depression as to OCD (Wu & Watson, 2005). In addition, studies of treatment outcome by symptom subtype have largely shown hoarding symptoms to predict poor outcome for standard OCD treatments using medication and exposure with response prevention (Mataix-Cols *et al.* 2002; Abramowitz *et al.* 2003; Steketee & Frost, 2003), suggesting that compulsive hoarding and OCD may involve different biological, cognitive or behavioral mechanisms.

* Address for correspondence: D. F. Tolin, Ph.D., Anxiety Disorders Center, The Institute of Living/Hartford Hospital, 200 Retreat Avenue, Hartford, CT 06106, USA.
(Email: dtolin@harthosp.org)

Hoarding behavior is thought to result in part from problems in decision making (Frost & Hartl, 1996; Steketee & Frost, 2003; Frost & Tolin, 2008). Research using the Iowa Gambling Task (Bechara *et al.* 1997), which requires participants to sacrifice immediate rewards to maximize long-term gain, shows that OCD patients with hoarding symptoms perform more poorly than do OCD patients without hoarding symptoms (Lawrence *et al.* 2006). This result was not replicated in a separate study (Grisham *et al.* 2007), although hoarding participants were noted to exhibit problems of impulsivity and inattention on standard neuropsychological tests. One possible explanation for the discrepancy between these two studies is the fact that Lawrence *et al.* (2006) included only OCD patients (with and without hoarding symptoms) whereas Grisham *et al.* (2007) sampled primary hoarding patients, 43% of who reported no other symptoms of OCD.

When asked to sort their possessions into categories, individuals with hoarding problems took more time, reported more anxiety and created a greater number of categories than did OCD patients and healthy controls (Wincze *et al.* 2007). Similar results were obtained using non-clinical volunteers who identified themselves as 'packrats' (Luchian *et al.* 2007). Importantly, however, hoarding participants in the Wincze *et al.* study showed no difficulty sorting objects that did not belong to them. Problems of executive functioning have also been noted among individuals with kleptomania (Grant *et al.* 2007), a condition that may overlap with hoarding in some respects.

Further complicating the decision-making process is the fact that individuals who hoard display maladaptive beliefs about their possessions, such as an exaggerated sense of emotional attachment to objects, memory-related concerns, responsibility for possessions, and desire for control over possessions (Steketee *et al.* 2003), as well as perfectionism and intolerance of uncertainty (Tolin *et al.* 2008). These factors are thought to render the decision-making process aversive and cumbersome, resulting in behavioral and cognitive avoidance (Frost *et al.* 1998, 2000).

The phenomenological and behavioral data described above suggest possible abnormalities in frontal cortical regions and anterior cingulate cortex (ACC). Impaired functioning of prefrontal cortex (PFC) and orbitofrontal cortex (OFC), for example, could be associated with diminished capacity to plan, organize, inhibit impulses, and anticipate behavioral consequences (Bechara *et al.* 1994; Rolls, 2004), as well as the diminished insight and self-awareness noted among individuals who hoard (Fitch *et al.* 2007). ACC abnormalities might account for the observed

problems in motivation, executive control, focused attention and misdirected emotional attachment seen in hoarding patients (Devinsky *et al.* 1995), as well as perfectionism, to the extent that this can be conceptualized as an erroneous sense of decisions being 'wrong' (Carter *et al.* 1998; Kiehl *et al.* 2000; Maltby *et al.* 2005).

Neuroimaging studies to date have broadly been consistent with the hypothesis that compulsive hoarding is associated with frontal cortical and ACC abnormality. Animal models of hoarding have implicated ACC (de Brabander *et al.* 1991). Three studies point to the neural underpinnings of compulsive hoarding in humans. In the first of these, resting positron emission tomography (PET) showed lower glucose metabolism in posterior cingulate and cuneus among hoarding patients ($n=12$) compared to healthy controls ($n=17$), and lower glucose metabolism in dorsal ACC compared to OCD patients ($n=33$) (Saxena *et al.* 2004). In the second study, functional magnetic resonance imaging (fMRI) was used with OCD patients ($n=16$) and healthy controls ($n=17$) during washing-, checking- and hoarding-related imaginal symptom provocation. For the hoarding provocation, participants were shown pictures of items, and were asked to imagine that they owned the items and would have to throw them away. During this task, OCD patients showed greater hemodynamic activity in left precentral gyrus and right medial OFC than did controls (Mataix-Cols *et al.* 2004). It is unclear, however, how these results would extend to compulsive hoarding populations because the study assessed the effects of viewing hoarding-related stimuli in patients with OCD, most of whom did not report high levels of compulsive hoarding symptoms. Finally, MRI was used with 63 adult patients with focal brain lesions. Nine of these patients reported the onset of excessive and impairing collecting and saving behavior (that topographically seems to resemble compulsive hoarding) following the onset of the neurological condition. Lesion locations among these nine patients differed from those of the 54 lesion patients who did not show such behaviors mainly in right mesial PFC, extending to ACC (Anderson *et al.* 2005).

The aim of the present pilot study was to explore whether the neural regions of interest derived from previous research (Mataix-Cols *et al.* 2004; Saxena *et al.* 2004) differentiate hoarding and non-hoarding participants during a decision-making task in which participants make actual, real-time decisions about whether to keep or discard their possessions. It was predicted that when making decisions about their own possessions, hoarding participants would show increased hemodynamic activity in OFC, PFC and

Table 1. Demographic and symptom information

Variable	Hoarding patients	Healthy controls
Age, mean (s.d.)	46.5 (10.10)	46.6 (10.55)
Female, <i>n</i> (%)	9 (75)	9 (75)
Caucasian, <i>n</i> (%)	11 (92)	11 (92)
Years of education, mean (s.d.)	16.46 (3.41)	16.44 (1.42)
HAMD total, mean (s.d.)	10.80 (4.16)	0.91(1.04)
SI-R total score, mean (s.d.)	50.07 (13.01)	10.0 (6.20)
Co-morbid Axis I diagnosis, <i>n</i> (%)	8 (67)	0 (0)
Major depressive disorder, <i>n</i> (%)	4 (33)	0 (0)
Obsessive-compulsive disorder, <i>n</i> (%)	2 (17)	0 (0)
Other anxiety disorder, <i>n</i> (%)	4 (33)	0 (0)
Co-morbid Axis II diagnosis, <i>n</i> (%)	4 (33)	0 (0)
Obsessive-compulsive disorder, <i>n</i> (%)	3 (25)	0 (0)
Other personality disorder, <i>n</i> (%)	3 (25)	0 (0)
Taking psychiatric medications, <i>n</i> (%)	8 (67)	0 (0)

HAMD, Hamilton Rating Scale for Depression; SI-R, Saving Inventory – Revised; s.d., standard deviation.

Percentages of co-morbid conditions sum to greater than 100% because some participants were diagnosed with multiple conditions.

ACC compared to control participants. No such differences were predicted when making decisions about control items that participants did not own, given the apparent absence of decision-making impairments for non-owned items (Wincze *et al.* 2007). Because of the exploratory nature of this study, whole-brain analyses were also conducted to generate hypotheses for future research.

Method

Participants

Twelve adult patients with compulsive hoarding and 12 matched healthy controls participated in the study and provided written informed consent. Table 1 presents demographic and clinical characteristics of the sample. All assessments were conducted by a doctoral-level psychologist with experience in the evaluation of compulsive hoarding, OCD and anxiety disorders. Consistent with previous research (Saxena *et al.* 2004), participants were classified as having compulsive hoarding if they met the clinical criteria for compulsive hoarding outlined by Frost & Hartl (1996), hoarding was their primary diagnosis as assessed by the evaluator, and the clinician's global impairment rating was 'moderately ill' or above. Where there were questions about the severity of compulsive hoarding, symptom severity was confirmed by a home visit or analysis of current photographs of living space. Compulsive hoarding

participants were excluded if they had a history of psychotic disorder, neurologic disorder, substance abuse, or serious suicidal ideation, or if compulsive hoarding was not their primary diagnosis. Concurrent major depressive disorder was allowed as long as it was not the primary diagnosis, given the high level of co-morbidity between hoarding and depression (Frost *et al.* 2006). Four of the compulsive hoarding participants were diagnosed with major depressive disorder, two with OCD, and four with other anxiety disorders; four also met criteria for a personality disorder. Most of the sample (67%) were taking psychiatric medications [selective serotonin (SSRIs) or serotonin norepinephrine (SNRIs) reuptake inhibitors]. None of the participants were taking high-potency benzodiazepines. Healthy controls were excluded if they met criteria for a current or past Axis I or Axis II disorder, had a history of neurological disorders, or were taking psychiatric medications.

Measures

Diagnostic status was determined using the Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-I; First *et al.* 1995) and the Structured Clinical Interview for DSM-IV Personality Disorders (SCID-II; First *et al.* 1997). Severity of compulsive hoarding symptoms was established using the Saving Inventory – Revised (SI-R; Frost *et al.* 2004). Symptom criteria for compulsive hoarding were established using the symptom checklist of the Yale-Brown

Obsessive–Compulsive Scale (YBOCS; Goodman *et al.* 1989) as well as the Hoarding Rating Scale – Interview (HRS-I; Tolin *et al.*, unpublished observations), which contains interviewer ratings of clutter, difficulty discarding, and acquisition from 0 (none) to 8 (extreme). Severity of depression was assessed using the Hamilton Rating Scale for Depression (HAM-D; Hamilton, 1960). Global impressions of illness severity were recorded using the Clinician’s Global Impression (CGI; Guy, 1976) scale. Subjective anxiety during the fMRI task was rated by participants on a scale from 0 (none) to 3 (extreme) by button-press.

Materials

The discarding task was designed to assess hemodynamic activity associated with successful and unsuccessful decisions to discard personal possessions and control items. For this task, we selected three different sets of stimuli to be discarded and destroyed:

- (1) Paper items (e.g. junk mail, newspapers) that belonged to the participants. Because compulsive hoarding participants have difficulty discarding ordinary possessions, we matched these items by type (e.g. junk mail, newspapers) rather than by emotional valence. Thus, compulsive hoarding participants were instructed to bring to the scanner session paper items meeting these criteria that they would normally have difficulty discarding. Healthy controls were instructed to bring to the scanner session the same types of paper items but without the restriction that they have difficulty discarding the items. We refer to these items as Participant’s Possessions (PP). These items were placed in a clear plastic bin on the right or left (counterbalanced) visual field. The bin was labeled with the participant’s first name.
- (2) Comparable paper items that did not belong to the participant. The research team saved their junk mail and old newspapers for a period of approximately 6 months. For each participant, we selected items that were roughly the same amount, size and type as the participant’s items. We refer to these items as Experimenter’s Possessions (EP). These items were placed in a separate clear plastic bin on the opposite side of the visual field from the PP. This bin was labeled with the experimenter’s first name.
- (3) Because of the possibility that participants would decline to discard a PP or an EP, we wanted to insure that any observed differences in hemodynamic response were not simply due to the visual stimulus of seeing an item being destroyed. Therefore, next to the PP and EP bins, we placed two clear plastic bins containing red pieces of



Fig. 1. Apparatus as seen from the participant’s perspective while in the scanner. In this picture, the experimenter is holding up a Participant’s Possession (PP, right hand) and a Neutral Stimulus (NS, left hand). Experimenter’s Possessions (EP) and their accompanying NS are on the experimenter’s left. The shredder is in the center of the picture.

blank paper that would be discarded and destroyed if the participant declined to discard a PP or an EP item. We refer to these items as Neutral Stimuli (NS).

The four clear plastic bins (PP, NS, EP, NS) were arranged on a table in front of a plain background, with the left/right location of the PP and EP bins counterbalanced. In the middle of a table was placed a document shredder that had been modified by removing the front cover so that items placed in the shredder could be seen clearly as they were being destroyed. A video camera was placed in front of the materials. Once in the scanner, participants viewed a live video feed of the apparatus that was rear-projected to an opaque screen mounted at the top of the bore of the magnet. Fig. 1 presents the layout of the materials as seen from the participant’s perspective while in the scanner.

Apparatus

Imaging was implemented on a Siemens 3T Allegra scanner. The Allegra is a high-performance head-dedicated system optimized for functional brain imaging. Head motion was restricted using a custom-built apparatus that interfaced with the head coil. Functional image volumes were collected with an echo planar imaging (EPI) gradient-echo pulse sequence [TR/TE 1500/28 ms, flip angle 65°, field of view (FOV) 24 × 24 cm, 64 × 64 matrix, 3.4 × 3.4 mm in plane resolution, 5 mm slice thickness, 29 slices] that effectively covers the entire brain (145 mm) in 1.5 s. The thin slices and short TE make the Allegra optimal for examining orbitofrontal activity.

Procedure

All study procedures were approved by the Institutional Review Board at Hartford Hospital. Participants gave written informed consent prior to the study. After being placed in the scanner and being given sufficient time to habituate to the environment, participants were alternately presented with PP/NS or EP/NS pairs through a live video feed and asked to make the decision of whether to discard the PP and EP items. The experimenter's presentation of each PP/NS or EP/NS pair served as the participant's cue to make the decision to shred that item. Participants were given 8 s to make this decision and to indicate their decision (keep *versus* shred) by button-press. The experimenter was cued by computer to either place the PP/EP item in the shredder (upon button-press) or to place the PP/EP item to the side and instead place the paired NS in the shredder (at 8 s, if the participant did not press the discard button). Once placed in the shredder, the items were visibly destroyed. Participants completed six blocks for each PP/NS or EP/NS pair. Each block lasted 24 s and consisted of three repetitions of each PP/NS or EP/NS pair.

Image processing

Functional images were reconstructed offline and re-oriented to approximately the anterior commissure/posterior commissure (AC/PC) plane. The two functional image runs were realigned using INRIAlign, a motion correction algorithm unbiased by local signal changes (Freire & Mangin, 2001; Freire *et al.* 2002), spatially normalized using custom linear and non-linear algorithms (Friston *et al.* 1995), smoothed, and analyzed in standard MNI space using Statistical Parametric Mapping (SPM2). Event-related responses were modeled using a synthetic hemodynamic response function composed of a single gamma function that modeled the hemodynamic response using a peak latency of 6 s. The modeled hemodynamic response for each run was derived by constructing a sequence of appropriately located synthetic responses for stimulus of interest. A 256-Hz high-pass filter was incorporated into the model to remove noise associated with low-frequency confounds (e.g. respiratory artifact). All images were normalized to a mean of 100 (arbitrary units) for each run to compensate for any intensity variations across runs.

For both groups, contrasts were specified on an individual subject basis that evaluated the effects of deciding to discard PP items relative to the baseline of discarding EP items, using images acquired during the 8-s decision-making period. Second-order movement parameters, the square of those generated from INRIAlign, were regressed out for each individual

Table 2. Reaction times, subjective anxiety ratings, and percentage discarded when deciding whether to discard Participant's Possessions (PP) and Experimenter's Possessions (EP)

Variable	Hoarding patients	Healthy controls
Reaction times (ms)		
PP	4531.88 (742.48) ^{a,b}	1960.76 (1020.56)
EP	2040.18 (932.30)	1656.69 (324.25)
Anxiety ratings, 0–3		
PP	0.98 (0.56) ^{a,b}	0.12 (0.17)
EP	0.45 (0.42) ^a	0.06 (0.15)
Percentage discarded		
PP	68.8 (19.0) ^{c,d}	97.7 (5.1)
EP	94.3 (5.6) ^d	98.6 (3.2)

^a Hoarding patients > healthy controls, $p < 0.05$.

^b PP > EP, $p < 0.05$.

^c EP > PP, $p < 0.05$.

^d Hoarding patients < healthy controls, $p < 0.05$.

subject contrast. The images containing these amplitudes were then entered into the second-level analyses (i.e. random effects analyses). Between-group comparisons were also entered into analyses of covariance (ANCOVAs) for compulsive hoarding participants having greater activation than healthy controls and for controls having greater activation than hoarding participants, both with HAMD scores as a covariate. Because of the exploratory nature of the present study, for all analyses a statistical threshold of $p < 0.01$ with eight contiguous voxels was used, and small volume correction (SVC) for the regions of interest found in previous research (Mataix-Cols *et al.* 2004; Saxena *et al.* 2004) was not conducted.

Results

Behavioral data

Table 2 presents reaction times during successful decisions to discard PP and EP items, subjective anxiety during the discarding task, and the percentage of refusals to discard PP and EP items for each group. As can be seen, compulsive hoarding participants responded differentially to PP *versus* EP items, whereas healthy controls did not. A 2 (group: hoarder, healthy) \times 2 (possession type: PP, EP) mixed-factor analysis of variance (ANOVA) yielded a significant main effect of group [$F(1, 19) = 41.81$, $p < 0.001$], a significant main effect of possession type [$F(1, 19) = 28.06$, $p < 0.001$], and a significant interaction [$F(1, 19) = 17.18$, $p = 0.001$]. Follow-up within-group analyses indicated that compulsive hoarding participants took significantly longer to decide to discard PP items than EP

items ($t_9=5.87$, $p<0.001$); there was no such difference for healthy controls ($t_{10}=0.94$, $p=0.369$). Comparing the two groups, compulsive hoarding participants took significantly longer than did healthy controls in deciding whether to discard PP items ($t_{19}=6.54$, $p<0.001$) but did not differ in the time it took them to decide whether to discard EP items ($t_{19}=1.28$, $p=0.214$).

Next, a 2 (group: hoarder, healthy) \times 2 (possession type: PP, EP) mixed-factor ANOVA of mean anxiety ratings (0–3) during the decision-making task yielded a significant main effect of group [$F(1, 19)=22.30$, $p<0.001$], a significant main effect of possession type [$F(1, 19)=12.82$, $p=0.002$], and a significant interaction [$F(1, 19)=8.05$, $p=0.011$]. Follow-up between-group analyses indicated that compulsive hoarding participants reported significantly greater anxiety than did healthy controls when deciding to discard both PP items ($t_{19}=4.90$, $p<0.001$) and EP items ($t_{19}=2.92$, $p=0.009$). Hoarding participants reported greater subjective anxiety when making decisions about whether to discard PP items than EP items ($t_9=3.20$, $p=0.011$). No difference in anxiety ratings between PP and EP items was observed for healthy controls ($t_{10}=1.29$, $p=0.226$). Pearson correlations of behavioral data with the HRS-I in hoarding participants indicated that subjective anxiety about discarding EP items was more strongly associated with acquisition ($r=0.30$) than with clutter ($r=0.15$) or difficulty discarding ($r=0.25$), although this correlation was not significant.

Finally, a 2 (group: hoarder, healthy) \times 2 (possession type: PP, EP) mixed-factor ANOVA of rates of discarding *versus* keeping items yielded a significant main effect of group [$F(1, 19)=25.59$, $p<0.001$], a significant main effect of possession type [$F(1, 19)=16.63$, $p=0.001$], and a significant interaction [$F(1, 19)=14.63$, $p=0.001$]. Follow-up within-group analyses indicated that compulsive hoarding participants discarded significantly fewer PP items than EP items ($t_9=3.98$, $p=0.003$); there was no such difference for healthy controls ($t_{10}=0.41$, $p=0.692$). Comparing the two groups, compulsive hoarding participants discarded significantly fewer PP items than did healthy controls ($t_{19}=4.86$, $p<0.001$); they also discarded significantly fewer EP items than did healthy controls ($t_{19}=2.15$, $p=0.045$). Pearson correlations indicated that the percentage of PP items discarded was more strongly associated with HRS-I difficulty discarding ($r=-0.54$) than with clutter ($r=0.06$) or acquisition ($r=-0.01$) in the hoarding participants. Thus, the behavioral data suggest that the discarding task was a successful analogue of hoarding behaviors in that compulsive hoarding participants had selective difficulty discarding their own possessions whereas healthy controls did not.




Fig. 2. Areas of significantly decreased hemodynamic activity for compulsive hoarding participants (□) *versus* healthy controls (○) while deciding whether to discard Participant's Possessions (PP) *versus* baseline. Image is thresholded at $p<0.01$.

fMRI results

Because compulsive hoarding is associated with depression (Frost *et al.* 2000), we used an ANCOVA, with HAMD scores as the covariate, for between-group analyses of the hemodynamic response. Examining activation during PP decisions *versus* baseline (see Fig. 2), hoarding participants did not show greater activity than did healthy controls in any regions. However, healthy controls showed greater hemodynamic activity than did hoarding participants in the left superior frontal gyrus ($-28, -4, 68$) and the left precentral gyrus (the latter region possibly reflecting the decreased button-pressing in this condition as described in *Behavioral data* above) [$-28, -16, 72$ (Brodmann 6)].

Next, we examined activation during PP decisions *versus* EP decisions (see Fig. 3). Group differences in the hemodynamic response were confined to the left hemisphere. Compulsive hoarding participants





Fig. 3. Areas of significantly increased hemodynamic activity for compulsive hoarding participants (\square) versus healthy controls (\square) while deciding whether to discard Participant's Possessions (PP) versus Experimenter's Possessions (EP). Image is thresholded at $p < 0.01$.

exhibited excessive activation in left lateral OFC [MNI coordinates $-36, 28, -20$ (Brodmann 47)], left amygdala and parahippocampal gyrus extending into thalamus ($-12, -8, -12$), and left cerebellum ($-8, -72, -20$) ($p < 0.01$) when compared to healthy controls. Healthy controls exhibited greater hemodynamic response than did compulsive hoarding participants only in the right lingual gyrus ($20, -84, 4$) ($p < 0.01$).

We also analyzed events where compulsive hoarding participants refused to discard PP items. This analysis was limited to compulsive hoarding participants because healthy controls did not refuse to discard their own possessions at rates that could be analyzed. For this contrast, we examined the hemodynamic response during events where compulsive hoarding participants refused to discard PP items when compared to successful decisions to discard PP items, again controlling for HAMD scores. Refusal to discard PP items (see Fig. 4) was associated with greater hemodynamic activity in superior temporal gyrus [$-40, 16, -40$ (Brodmann 38)], middle temporal gyrus ($-56, -4, -16$), medial frontal gyrus




Fig. 4. Areas of significant activation for compulsive hoarding participants when refusing to discard Participant's Possessions (PP) versus successful decisions to discard PP. Image is thresholded at $p < 0.01$.

[$4, 40, -16$ (Brodmann 11)], anterior cingulate extending to OFC ($-20, 32, 24$), precentral gyrus [$60, 0, 8$ (Brodmann 6)], and cerebellum ($8, -36, -28$; $16, -80, 20$) ($p < 0.01$); the latter finding was largely inside the ventricle and therefore probably an artifact. Non-significant activation was also seen in the cuneus.

Correlations with measures of symptom severity for compulsive hoarding participants

Pearson product moment correlations were calculated between compulsive hoarding symptom severity, as measured by the SI-R, and the hemodynamic response during decisions to discard PP items and refusals to discard PP items, controlling for depression as measured by the HAMD. During successful decisions to discard PP items, compulsive hoarding symptom severity was positively associated with hemodynamic response in subcallosal gyrus ($8, 12, -12$) [$8, 4, -16$ (Brodmann 34)], ($16, 8, -16$) and negatively associated with hemodynamic response in right inferior temporal gyrus [$52, -12, -36$ (Brodmann 20)], left middle temporal gyrus [$-52, -4, -24$ (Brodmann 21)], and right precentral gyrus ($56, 0, 48$). We examined whether these regions (and the 10 mm regions surrounding them) differentiated hoarding from healthy participants; only inferior temporal gyrus was less active in hoarding participants than in healthy controls.

During refusals to discard PP items, compulsive hoarding symptom severity was positively associated with hemodynamic response in globus pallidus ($-16, 0, -8$), lingual gyrus ($16, -64, -12$), and left cerebellum ($-36, -64, -16$). There were no areas of negative association for this contrast. We examined




Fig. 5. Scatterplots of correlations between hoarding severity (Saving Inventory – Revised, SI-R) and neural regions of interest during (a) successful decisions to discard Participant’s Possessions (PP) and (b) refusals to discard PP.

whether these regions (and the 10 mm regions surrounding them) differentiated hoarding from healthy participants; globus pallidus and lingual gyrus (but not cerebellum) showed greater activity in hoarding *versus* healthy participants. Scatterplots for these relationships are shown in Fig. 5.

Discussion

We found excessive hemodynamic activity in lateral OFC, a region that is associated with processing relative reward value, particularly values that are

experienced as punishing (Kringelbach & Rolls, 2004; Kringelbach, 2005). Increased OFC activity may also be associated with greater emotional engagement in affective decision making (Northoff *et al.* 2006). This finding is broadly consistent with the theory that hoarding behavior stems, in part, from abnormalities in the decision-making process (Frost & Hartl, 1996; Steketee & Frost, 2003; Frost & Tolin, 2008); the finding of increased hemodynamic response in lateral OFC may reflect hoarding patients’ greater sense that discarding low absolute worth personal possessions is punishing. It remains unclear whether hoarding

patients show a general deficit in decision-making capacity, or whether decision-making impairment is limited to personally relevant possessions. As described previously, research on this topic has yielded mixed results (Lawrence *et al.* 2006; Grisham *et al.* 2007), and more research is needed.

Patterns of hemodynamic activity observed in healthy controls are also generally consistent with those from a healthy volunteer study in which participants imagined discarding possessions and exhibited hemodynamic activity in prefrontal regions that did not differ from activity associated with normally aversive tasks (Mataix-Cols *et al.* 2003).

One potential limitation of the present study is the fact that decisions about whether to keep *versus* discard items were interspersed with the shredding of items, raising the possibility that the neural mechanisms of decision making were confounded by an aversive response to seeing items discarded and shredded. Although images were acquired during the 8-s decision-making process, because of the sequence of events (decide, discard, decide, discard, etc.) it is possible that neural activity during decision making could include some carryover effects of the discarding and shredding procedure. Future studies should increase the temporal separation of these two events to clarify whether the obtained results are specific to the decision-making process. It would also be helpful for future research to compare neural decision-making processes with neuropsychological indices of impaired decision making.

Also consistent with the notion of difficulty in the decision-making process is the finding of greater activity in left parahippocampal gyrus in hoarding *versus* control participants. Activity in this region is also associated with effortful memory search and retrieval (Gur *et al.* 1997; Maguire & Mummery, 1999), including memory for the associative properties of objects (Yonelinas *et al.* 2001). The left parahippocampal gyrus has also been associated with the processing of unpleasant emotions (Lane *et al.* 1997); the difference between groups does not seem to be solely attributable to depression, which was covaried in the present analyses. Relatedly (although not expected), when examining activation during PP decisions *versus* baseline (rather than EP decisions), hoarding participants showed reduced activity in left superior frontal gyrus compared to healthy controls. The left superior frontal gyrus has been associated with working memory, particularly with more challenging tasks (du Boisgueheneuc *et al.* 2006); suppression of activity in superior frontal gyrus may reflect impaired capacity for self-reflection due to cognitive demand (Goldberg *et al.* 2006).

Refusals to discard possessions might provide the closest approximation to actual hoarding behaviors. The strength of conclusions that can be drawn from this analysis is limited by the fact that healthy controls refused to discard only a very small number of items, precluding between-group comparisons. For compulsive hoarding participants, refusal to discard personal possessions was associated with increased hemodynamic response, compared to control items, in middle frontal gyrus extending into rostral ACC, a region associated with effortful learning and problem solving, error detection, motivation, and modulation of emotional responses (Devinsky *et al.* 1995; Carter *et al.* 1998; Bush *et al.* 2000; Kiehl *et al.* 2000; Allman *et al.* 2001; Polli *et al.* 2005). Thus, ACC abnormalities might reflect the excessive emotional attachment and exaggerated concerns about making the wrong decision reported by individuals with compulsive hoarding. It is noted that hoarding participants refused to discard only 31% of their personal possessions, raising questions of generalizability to actual hoarding situations. One possibility is that participants pre-sorted their possessions prior to the experiment, and did not bring in the most challenging items. Another possibility is that demand characteristics of the experiment increased participants' willingness to discard.

The present results differ from those obtained by Saxena *et al.* (2004), who observed lower glucose metabolism in posterior cingulate and cuneus when compared to healthy controls, and in dorsal ACC when compared to non-hoarding OCD patients. That study examined neural activity at rest, rather than during a decision-making challenge. One possible explanation for the discrepancy is a biphasic pattern of neural activity in compulsive hoarding, with decreased activity at rest but increased activity during decision making. Additional research is needed to clarify the time course of neural activity in compulsive hoarding. Mataix-Cols *et al.* (2004) found greater hemodynamic activity in left precentral gyrus and right medial OFC when participants with OCD (most of whom did not appear to suffer from significant hoarding symptoms) were asked to imagine discarding objects. The present finding of increased OFC and precentral activity during actual decision making might be consistent with those results, although the location of activity differed. Our results are also broadly consistent with animal models of hoarding suggesting the role of ACC (de Brabander *et al.* 1991) as well as hippocampus (Kolb, 1977) and globus pallidus (Mogenson & Wu, 1988) in food-hoarding behavior.

Much of the significant hemodynamic activity in the present study was found in association cortex (e.g.

parahippocampal regions, lingual gyrus), particularly temporal lobe (superior and middle temporal gyrus), and motor regions (e.g. cerebellum, precentral gyrus). These were unexpected findings and await clarification in a larger sample, although the motor findings might have been a methodological artifact; because participants pressed a button to signal an intent to discard an item, and refrained from pressing a button to signal an intent to keep an item, different responses might well have differentially recruited motor regions of the brain. Future work should use button-press for intent to keep, as well as discard, an item in order to rule out this potential confound.

The present findings may aid understanding of the refractory nature of compulsive hoarding. Individuals with this condition may find the act of deciding to discard personal possessions cognitively burdensome, punishing and aversive, and hence are likely to avoid such decisions in the future. Avoidance of decision making, in turn, leads to the continued accumulation of clutter, ultimately resulting in substantial functional impairment. Because the present results cannot distinguish the effects of compulsive hoarding from those of psychopathology more broadly (and it is noted again that the majority of the sample met criteria for at least one co-morbid psychiatric condition), future studies should use clinical control groups to differentiate the hemodynamic responses of hoarding patients from those of patients with other clinical conditions such as OCD. As only a minority of the present participants met criteria for non-hoarding OCD, it remains unclear whether individuals suffering from co-morbid OCD and hoarding would show similar patterns of neural activity.

Acknowledgments

This work was supported by Hartford Hospital grant 126083 (to N.M.) and NIMH grant R01 MH074934 (to D.F.T.).

Declaration of Interest

None.

References

- Abramowitz JS, Franklin ME, Schwartz SA, Furr JM (2003). Symptom presentation and outcome of cognitive-behavioral therapy for obsessive-compulsive disorder. *Journal of Consulting and Clinical Psychology* **71**, 1049–1057.
- Allman JM, Hakeem A, Erwin JM, Nimchinsky E, Hof P (2001). The anterior cingulate cortex. The evolution of an interface between emotion and cognition. *Annals of the New York Academy of Sciences* **935**, 107–117.
- Anderson SW, Damasio H, Damasio AR (2005). A neural basis for collecting behaviour in humans. *Brain* **128**, 201–212.
- Bechara A, Damasio AR, Damasio H, Anderson SW (1994). Insensitivity to future consequences following damage to human prefrontal cortex. *Cognition* **50**, 7–15.
- Bechara A, Damasio H, Tranel D, Damasio AR (1997). Deciding advantageously before knowing the advantageous strategy. *Science* **275**, 1293–1295.
- Bush G, Luu P, Posner MI (2000). Cognitive and emotional influences in anterior cingulate cortex. *Trends in Cognitive Science* **4**, 215–222.
- Calamari JE, Wiegartz PS, Riemann BC, Cohen RJ, Greer A, Jacobi DM, Jahn SC, Carmin C (2004). Obsessive-compulsive disorder subtypes: an attempted replication and extension of a symptom-based taxonomy. *Behaviour Research and Therapy* **42**, 647–670.
- Carter CS, Braver TS, Barch DM, Botvinick M, Noll D, Cohen J (1998). Anterior cingulate cortex, error detection, and the online monitoring of performance. *Science* **280**, 747–749.
- de Brabander JM, de Bruin JP, van Eden CG (1991). Comparison of the effects of neonatal and adult medial prefrontal cortex lesions on food hoarding and spatial delayed alternation. *Behavioral Brain Research* **42**, 67–75.
- Devinsky O, Morrell MJ, Vogt BA (1995). Contributions of anterior cingulate cortex to behaviour. *Brain* **118**, 279–306.
- du Boisgueheneuc F, Levy R, Volle E, Seassau M, Duffau H, Kinkingnehun S, Samson Y, Zhang S, Dubois B (2006). Functions of the left superior frontal gyrus in humans: a lesion study. *Brain* **129**, 3315–3328.
- First MB, Spitzer RL, Gibbon M, Williams JBW (1995). *Structured Clinical Interview for DSM-IV Axis I Disorders – Patient Edition (SCID-I/P, version 2.0)*. Biometrics Research Department: New York.
- First MB, Spitzer RL, Gibbon M, Williams JBW (1997). *Structured Clinical Interview for DSM-IV Personality Disorders (SCID-II)*. American Psychiatric Press, Inc.: Washington, DC.
- Fitch KE, Tolin DF, Frost RO, Steketee G (2007). Compulsive hoarding: assessing insight. Presented at the Annual Meeting of the Association of Behavioral and Cognitive Therapies, Philadelphia, November, 2007.
- Freire L, Mangin JF (2001). Motion correction algorithms may create spurious brain activations in the absence of subject motion. *NeuroImage* **14**, 709–722.
- Freire L, Roche A, Mangin JF (2002). What is the best similarity measure for motion correction in fMRI time series? *IEEE Transactions on Medical Imaging* **21**, 470–484.
- Friston KJ, Ashburner J, Frith CD, Poline J-B, Heather JD, Frackowiak RSJ (1995). Spatial registration and normalization of images. *Human Brain Mapping* **2**, 165–189.
- Frost RO, Hartl TL (1996). A cognitive-behavioral model of compulsive hoarding. *Behaviour Research and Therapy* **34**, 341–350.
- Frost RO, Kim HJ, Morris C, Bloss C, Murray-Close M, Steketee G (1998). Hoarding, compulsive buying and

- reasons for saving. *Behaviour Research and Therapy* **36**, 657–664.
- Frost RO, Steketee G, Grisham J** (2004). Measurement of compulsive hoarding: Saving Inventory – Revised. *Behaviour Research and Therapy* **42**, 1163–1182.
- Frost RO, Steketee G, Tolin DF, Brown TA** (2006). Comorbidity and diagnostic issues in compulsive hoarding. Presented to the Annual Meeting of the Anxiety Disorders Association of America/>, Miami, Florida, March 2006.
- Frost RO, Steketee G, Williams LF, Warren R** (2000). Mood, personality disorder symptoms and disability in obsessive compulsive hoarders: a comparison with clinical and nonclinical controls. *Behaviour Research and Therapy* **38**, 1071–1081.
- Frost RO, Tolin DF** (2008). Compulsive hoarding. In *Clinical Handbook of Obsessive-Compulsive Disorder and Related Problems* (ed. S. Taylor, J. S. Abramowitz and D. McKay), pp. 76–94. Johns Hopkins University Press: Baltimore.
- Goldberg II, Harel M, Malach R** (2006). When the brain loses its self: prefrontal inactivation during sensorimotor processing. *Neuron* **50**, 329–339.
- Goodman WK, Price LH, Rasmussen SA, Mazure C, Fleischmann RL, Hill CL, Heninger GR, Charney DS** (1989). The Yale-Brown Obsessive Compulsive Scale. I. Development, use, and reliability. *Archives of General Psychiatry* **46**, 1006–1011.
- Grant JE, Odlaug BL, Wozniak JR** (2007). Neuro-psychological functioning in kleptomania. *Behaviour Research and Therapy* **45**, 1663–1670.
- Grisham JR, Brown TA, Savage CR, Steketee G, Barlow DH** (2007). Neuropsychological impairment associated with compulsive hoarding. *Behaviour Research and Therapy* **45**, 1471–1483.
- Gur RC, Ragland JD, Mozley LH, Mozley PD, Smith R, Alavi A, Bilker W, Gur RE** (1997). Lateralized changes in regional cerebral blood flow during performance of verbal and facial recognition tasks: correlations with performance and ‘effort’. *Brain and Cognition* **33**, 388–414.
- Guy W** (1976). *Assessment Manual for Psychopharmacology*. U.S. Government Printing Office: Washington, DC.
- Hamilton M** (1960). A rating scale for depression. *Journal of Neurological and Neurosurgical Psychiatry* **23**, 56–62.
- Kiehl KA, Liddle PF, Hopfinger JB** (2000). Error processing and the rostral anterior cingulate: an event-related fMRI study. *Psychophysiology* **37**, 216–223.
- Kolb B** (1977). Studies on the caudate-putamen and the dorsomedial thalamic nucleus of the rat: implications for mammalian frontal-lobe functions. *Physiology and Behavior* **18**, 237–244.
- Kringelbach ML** (2005). The human orbitofrontal cortex: linking reward to hedonic experience. *Nature Reviews Neuroscience* **6**, 691–702.
- Kringelbach ML, Rolls ET** (2004). The functional neuroanatomy of the human orbitofrontal cortex: evidence from neuroimaging and neuropsychology. *Progress in Neurobiology* **72**, 341–372.
- Lane RD, Reiman EM, Bradley MM, Lang PJ, Ahern GL, Davidson RJ, Schwartz GE** (1997). Neuroanatomical correlates of pleasant and unpleasant emotion. *Neuropsychologia* **35**, 1437–1444.
- Lawrence NS, Wooderson S, Mataix-Cols D, David R, Speckens A, Phillips ML** (2006). Decision making and set shifting impairments are associated with distinct symptom dimensions in obsessive-compulsive disorder. *Neuropsychology* **20**, 409–419.
- Luchian SA, McNally RJ, Hooley JM** (2007). Cognitive aspects of nonclinical obsessive-compulsive hoarding. *Behaviour Research and Therapy* **45**, 1657–1662.
- Maguire EA, Mummery CJ** (1999). Differential modulation of a common memory retrieval network revealed by positron emission tomography. *Hippocampus* **9**, 54–61.
- Maltby N, Tolin DF, Worhunsky P, O’Keefe TM, Kiehl KA** (2005). Dysfunctional action monitoring hyperactivates frontal-striatal circuits in obsessive-compulsive disorder: an event-related fMRI study. *NeuroImage* **24**, 495–503.
- Mataix-Cols D, Cullen S, Lange K, Zelaya F, Andrew C, Amaro E, Brammer MJ, Williams SC, Speckens A, Phillips ML** (2003). Neural correlates of anxiety associated with obsessive-compulsive symptom dimensions in normal volunteers. *Biological Psychiatry* **53**, 482–493.
- Mataix-Cols D, Marks IM, Greist JH, Kobak KA, Baer L** (2002). Obsessive-compulsive symptom dimensions as predictors of compliance with and response to behaviour therapy: results from a controlled trial. *Psychotherapy and Psychosomatics* **71**, 255–262.
- Mataix-Cols D, Wooderson S, Lawrence N, Brammer MJ, Speckens A, Phillips ML** (2004). Distinct neural correlates of washing, checking, and hoarding symptom dimensions in obsessive-compulsive disorder. *Archives of General Psychiatry* **61**, 564–576.
- Meunier SA, Tolin DF, Frost RO, Steketee G, Brady RE** (2006). Prevalence of hoarding symptoms across the anxiety disorders. Presented to the Annual Meeting of the Anxiety Disorders Association of America, Miami, FL, March 2006.
- Mogenson GJ, Wu M** (1988). Disruption of food hoarding by injections of procaine into mediodorsal thalamus, GABA into subpallidal region and haloperidol into accumbens. *Brain Research Bulletin* **20**, 247–251.
- Northoff G, Grimm S, Boeker H, Schmidt C, BERPohl F, Heinzel A, Hell D, Boesiger P** (2006). Affective judgment and beneficial decision making: ventromedial prefrontal activity correlates with performance in the Iowa Gambling Task. *Human Brain Mapping* **27**, 572–587.
- Polli FE, Barton JJ, Cain MS, Thakkar KN, Rauch SL, Manoach DS** (2005). Rostral and dorsal anterior cingulate cortex make dissociable contributions during antisaccade error commission. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 15700–15705.
- Rolls ET** (2004). The functions of the orbitofrontal cortex. *Brain and Cognition* **55**, 11–29.
- Saxena S, Brody AL, Maidment KM, Smith EC, Zohrabi N, Katz E, Baker SK, Baxter Jr LR** (2004). Cerebral glucose

- metabolism in obsessive-compulsive hoarding. *American Journal of Psychiatry* **161**, 1038–1048.
- Steketee G, Frost RO** (2003). Compulsive hoarding: current status of the research. *Clinical Psychology Review* **23**, 905–927.
- Steketee G, Frost RO, Kim HJ** (2001). Hoarding by elderly people. *Health and Social Work* **26**, 176–184.
- Steketee G, Frost RO, Kyrios M** (2003). Beliefs about possessions among compulsive hoarders. *Cognitive Therapy and Research* **27**, 467–479.
- Tolin DF, Brady RE, Hannan SE** (2008). Obsessional beliefs and symptoms of obsessive-compulsive disorder in a clinical sample. *Journal of Psychopathology and Behavioral Assessment* **30**, 31–42.
- Tolin DF, Frost RO, Steketee G, Gray KD, Fitch KE** (2007). The economic and social burden of compulsive hoarding. *Psychiatry Research*. doi: 10.1016/j.psychres.2007.08.008.
- Wincze JP, Steketee G, Frost RO** (2007). Categorization in compulsive hoarding. *Behaviour Research and Therapy* **45**, 63–72.
- Wu KD, Watson D** (2005). Hoarding and its relation to obsessive-compulsive disorder. *Behaviour Research and Therapy* **43**, 897–921.
- Yonelinas AP, Hopfinger JB, Buonocore MH, Kroll NE, Baynes K** (2001). Hippocampal, parahippocampal and occipital-temporal contributions to associative and item recognition memory: an fMRI study. *Neuroreport* **12**, 359–363.