

Epigenetic Status of *Gdnf* in the Ventral Striatum Determines Susceptibility and Adaptation to Daily Stressful Events

Shusaku Uchida,^{1,*} Kumiko Hara,¹ Ayumi Kobayashi,¹ Koji Otsuki,¹ Hirotaka Yamagata,¹ Teruyuki Hobara,¹ Takayoshi Suzuki,² Naoki Miyata,² and Yoshifumi Watanabe^{1,*}

¹Division of Neuropsychiatry, Department of Neuroscience, Yamaguchi University Graduate School of Medicine, 1-1-1 Minamikogushi, Ube, Yamaguchi 755-8505, Japan

²Graduate School of Pharmaceutical Sciences, Nagoya City University, 3-1 Tanabe-dori, Mizuho-ku, Nagoya, Aichi 467-8603, Japan

*Correspondence: s-uchida@yamaguchi-u.ac.jp (S.U.), yoshiwat@yamaguchi-u.ac.jp (Y.W.)

DOI 10.1016/j.neuron.2010.12.023

SUMMARY

Stressful events during adulthood are potent adverse environmental factors that can predispose individuals to psychiatric disorders, including depression; however, many individuals exposed to stressful events can adapt and function normally. While stress vulnerability may influence depression, the molecular mechanisms underlying the susceptibility and adaptation to chronic stress within the brain are poorly understood. In this study, two genetically distinct mouse strains that exhibit different behavioral responses to chronic stress were used to demonstrate how the differential epigenetic status of the glial cell-derived neurotrophic factor (*Gdnf*) gene in the ventral striatum modulates susceptibility and adaptation to chronic stress. Our results suggest that the histone modifications and DNA methylation of the *Gdnf* promoter have crucial roles in the control of behavioral responses to chronic stress. Our data provide insights into these mechanisms, suggesting that epigenetic modifications of *Gdnf*, along with genetic and environmental factors, contribute to behavioral responses to stress.

INTRODUCTION

Major depressive disorder is one of the most common and serious health problems in societies worldwide. While the etiology of this disorder is multifactorial and poorly understood, both genetic and environmental factors may be involved in the precipitation of depression (Charney and Manji, 2004; Krishnan and Nestler, 2008; Feder et al., 2009). Chronic stressful life events during adulthood are potent adverse environmental factors that can activate or amplify the expression of depression symptoms (Leonardo and Hen, 2008). Many individuals exposed to stressful events do not show signs or symptoms of depression; however, some individuals exposed to psychological stress are predisposed to major depression (Charney, 2004). Thus far, the

molecular mechanisms underlying the susceptibility and adaptation to chronic stress within the brain are poorly understood.

Genetically distinct mouse strains that exhibit substantial differences in anxiety and stress reactivity have been used as animal models for investigating the influence of genetic and environmental factors on brain functions and behaviors (Francis et al., 2003; Hovatta et al., 2005; Mozhui et al., 2010). In particular, the inbred BALB/c (BALB) mouse strain demonstrates unique responses to stress. Compared to the C57BL/6 (B6) stress-resilient strain, BALB mice show maladaptive responses to stressful stimuli (Francis et al., 2003; Hovatta et al., 2005; Bhansali et al., 2007; Palumbo et al., 2009). Therefore, BALB mice are considered a stress-vulnerable strain, and comparing the stress responses of BALB and B6 mice may provide useful information regarding the mechanisms of susceptibility and adaptation to stressful stimuli in brain function and behavior, such as those associated with depression.

Neuronal activity regulates a complex program of gene expression that is involved in the structural and functional plasticity of the brain (Flavell and Greenberg, 2008). There is also increasing evidence indicating that aberrant transcription regulation is one of the key components in the pathophysiology of depression (Tsankova et al., 2007; Krishnan and Nestler, 2008; Feder et al., 2009). Recent reports have suggested that the epigenetic regulation of genes, such as DNA methylation and histone modification, can trigger the development of stress vulnerability and contribute to the behavioral responses to chronic stress and antidepressants (Weaver et al., 2004; Tsankova et al., 2006; Fyffe et al., 2008; Jakobsson et al., 2008; LaPlant et al., 2010). However, the role of environmental factors along with genetic factors in the epigenetic regulation of the pathogenesis of depression is largely unknown.

The aim of the present study was to clarify the molecular mechanisms underlying the susceptibility and adaptation to chronic stress using stress-vulnerable BALB and stress-resilient B6 mice strains. Our results show that the differential epigenetic status of the glial cell-derived neurotrophic factor (*Gdnf*) gene in the nucleus accumbens (NAc) influences differential behavioral responses to stress. Therefore, we propose that epigenetic regulation of *Gdnf* by environmental factors, along with genetic factors, contributes to the level of susceptibility and adaptation ability of individuals to chronic stressful life events.

Table 1. Summary of Behavioral Characterizations of B6 and BALB Mice Subjected to CUMS

Behavioral Assay	Phenotype Tested	Stressed B6	Stressed BALB	Stressed BALB with IMI	Nonstressed BALB with IMI
		Versus Nonstressed B6	Versus Nonstressed BALB		
Forced swim test	Immobility time	↔	↑	↔	↓
	Latency to immobility	↔	↓	↔	↑
	First immobility time	↔	↑	↔	↔
Sucrose preference test	Preference ratio	↔	↓	↔	↔
	Total (water + sucrose) intake	↔	↔	↔	↔
Social interaction test	Interaction time	↑	↓	↔	↔
	Total number of interactions	↔	↓	↔	↔
Novelty-suppressed feeding test	Latency to feed	↓	↑	↔	↓
	Food consumption	↔	↔	↔	↔
	Body weight loss	↔	↔	↔	↔

This table shows the behavioral differences between B6 and BALB mice subjected to CUMS conditions for 6 weeks. Also shown are the effects of 3 weeks of antidepressant treatment in stressed and non-stressed BALB mice. ↔, no change; ↑, significantly greater changes; ↓, significantly fewer changes. IMI; imipramine.

RESULTS

Complete statistical summaries of behavior, gene expression by quantitative real-time PCR (Q-PCR) and Western blotting, and chromatin immunoprecipitation (ChIP) data are provided in Tables S1, S2, and S3 (available online), respectively.

Differential Behavioral Responses to Chronic Stress in B6 and BALB Mice

We first investigated the behavioral consequences of 6 weeks of chronic ultra-mild stress (CUMS) exposure, a procedure based solely on environmental and social stressors that do not include food or water deprivation (Lanfumey et al., 1999; Rangon et al., 2007), in BALB and B6 mice. The experimental design is shown in Figure S1A, and the results are summarized in Table 1. Anhedonia, diminished interest or pleasure, is one of the core symptoms of major depression (Wong and Licinio, 2001). Therefore, we examined whether this trait was present in stressed BALB mice using a sucrose preference test (Figures S2A and S2B). CUMS significantly decreased sucrose preference, and this effect was reversed by continuous treatment (via drinking water) with imipramine (IMI, 18 mg/kg/day), a tricyclic antidepressant (Figure S2A). Total fluid intake was not affected by either treatment (Figure S2B). We then subjected BALB mice to the acute forced swim test, which uses increased immobility time as an index of behavioral despair (Porsolt et al., 1977). CUMS significantly increased immobility times (Figure S2C) and the duration of the first immobility episode (Figure S2D) and reduced the latency to the first immobility episode (Figure S2E). These behavioral effects were reversed with continuous IMI treatment (Figures S2C–S2E).

Anxiety is frequently comorbid in patients with major depression. To examine the effects of CUMS on anxiety behavior, we performed the novelty-suppressed feeding test. The latency to begin eating in a novel environment has been used as an index of anxiety behavior (Richardson-Jones et al., 2010). Stressed BALB mice showed significantly longer latency periods to

feeding (Figure S2F), with no significant differences in weight loss induced by food deprivation (Figure S2G) or feeding activities (Figure S2H). Furthermore, the increased latency to feed induced by CUMS was reversed with continuous IMI treatment (Figure S2F). Anxiety behavior was also examined using the elevated zero maze test. The amount of time spent in the open section and frequency of rearing were not affected by CUMS (data not shown). Social interaction time also provides an index of anxiety and depression-like behavior. More anxious and depressed rodents spend less time in social interactions (File and Seth, 2003; Berton et al., 2006). Stressed BALB mice spent significantly less time engaged in social interactions and had fewer interactions than the nonstressed controls. This effect was also reversed with continuous IMI treatment (Figures S2I and S2J). Taken together, these results indicate an increase in depression- and anxiety-related behaviors in stressed BALB mice.

In contrast with the BALB mice, B6 mice subjected to CUMS did not show any behavioral changes in the sucrose preference test (Figures S3A and S3B) or forced swim test (Figures S3C and S3D), but they did demonstrate a reduced latency to feed in the novelty-suppressed feeding test (Figure S3E) and increased interaction times in the social interaction test (Figure S3G), suggesting a decrease in anxiety-related behaviors in stressed B6 mice. In addition to behavioral characterization, we also examined the plasma corticosterone (CORT) levels of mice to investigate how CUMS influences neuroendocrine function. We found increased plasma CORT levels 60 min after the initiation of a stressor in both BALB and B6 mice on day 3 of the CUMS session (Figures S4A and S4B). In contrast, on day 38 of the CUMS session, B6 mice showed a reduction in plasma CORT levels 60 min after the initiation of the stressor (Figure S4B). This effect was not observed in BALB mice (Figure S4A). Thus, BALB mice responded to CUMS with an increase in depression-like phenotypes, whereas the B6 mice responded to the same stress conditions with a decrease in anxiety-related behaviors. These behavioral and neuroendocrine data indicate




Figure 1. Differential Regulation of *Gdnf* Expression in Stress-Susceptible and Stress-Adaptive Mice Strains

(A) mRNA expression of *Gdnf* in the dSTR and vSTR of BALB mice subjected to CUMS or nonstress (NS) conditions with or without continuous IMI (18 mg/kg/day) treatment ($n = 6$ per group; * $p < 0.05$ versus NS mice receiving vehicle (normal water) in corresponding brain regions).

(B) mRNA expression of *Gdnf* in the dSTR and vSTR of B6 mice subjected to CUMS or NS conditions ($n = 6$ per group; * $p < 0.05$ versus NS in corresponding brain regions).

(C) Levels of GDNF proteins in the dSTR and vSTR of BALB and B6 mice subjected to CUMS or NS conditions with or without continuous IMI treatment ($n = 8-12$ per group; * $p < 0.05$ versus NS mice receiving water in corresponding brain regions).

(D–G) Correlation analyses of GDNF levels in the vSTR of nonstressed BALB (black line) and B6 (red line) mice and (D) the social interaction times (BALB; $n = 21$, B6; $n = 12$), (E) the sucrose preferences (BALB; $n = 28$, B6; $n = 16$), (F) the immobility times in the forced swim test (BALB; $n = 28$, B6; $n = 16$), and (G) the latency to feed in the novelty-suppressed feeding test (BALB; $n = 28$, B6; $n = 16$) (* $p < 0.05$).

(H and I) Successful transductions of EGFP (H) and GDNF (I) into the NAc using the PEI gene delivery system are shown.

(J and K) Effects of GDNF overexpression in the NAc of nonstressed B6 and stressed BALB mice ($n = 14-19$ per group) on social interaction times (J) and sucrose preference (K) (* $p < 0.05$ versus PEI/*Egfp* in corresponding strains). Data are presented as mean \pm SEM.

that BALB and B6 mice develop “passive” and “active” responses to stress, suggesting that these strains of mice are susceptible and adaptive strains to CUMS, respectively.

Expression Analyses of a Variety of Neurotrophic Factors in a Mouse Model of Depression

Neurotrophic factors play important roles in the regulation of synaptic and structural plasticity in the brain and may be involved in depression (Nestler et al., 2002; Duman and Monteggia, 2006). To investigate the contribution of neurotrophic factors to the behavioral abnormalities of stressed BALB mice, the mRNA levels of multiple neurotrophic factors were examined, including *Bdnf*, *Gdnf*, *Vegf*, *Nt-3*, *Nt-4/5*, *Cdnf*, *Ngf*, *Fgf2*, and *Igf1*, in regions of the brain associated with stress, such as the hippocampus (HP), prefrontal cortex, amygdala, striatum (STR), and hypothalamus, of BALB mice subjected to 6 weeks of CUMS either with or without continuous IMI treatment.

Q-PCR revealed that the expression levels of *Bdnf*, *Vegf*, and *Igf1* mRNA were significantly increased by continuous IMI treatment, but were not affected by CUMS (Figures S5B, S5D, and S5H). Interestingly, the mRNA levels of *Gdnf* and *Nt-3* in the STR and HP, respectively, were significantly decreased by CUMS, and these effects were reversed by continuous IMI treatment (Figures S5A and S5E). In addition, the mRNA expression level of *Gdnf* in stressed BALB mice was significantly decreased in both the dorsal STR (dSTR) and the ventral STR (vSTR) (Figure 1A). On the contrary, the mRNA expression level of *Gdnf* in stressed B6 mice was significantly increased in the vSTR but not in the dSTR (Figure 1B). These changes in GDNF expression were confirmed at the protein level using an ELISA assay (Figure 1C). These results suggest that the transcriptional regulation of *Gdnf* in the vSTR is differentially regulated in the two mouse strains and may contribute to the observed behavioral responses to CUMS.

Role of GDNF in the NAc in Depression-like Behaviors

We next investigated whether a correlation exists between *Gdnf* expression in the vSTR and behavioral performances in mice. We found that GDNF protein levels in the vSTR of nonstressed BALB and B6 mice were significantly correlated with social interaction time (Figure 1D) and sucrose preferences (Figure 1E), but not with immobility times in the forced swim test (Figure 1F) or the latency to feed in the novelty-suppressed feeding test (Figure 1G). These data suggest an important role for GDNF in the vSTR for determining certain types of depression-like behaviors.

To directly investigate the role of GDNF in depression-like behaviors, GDNF was overexpressed in the NAc of mice using the polyethylenimine (PEI) gene delivery system. The experimental design is shown in Figure S1B. The successful transduction of EGFP (Figure 1H) and GDNF (Figure 1I) into the NAc of mice using this system was confirmed. We first assessed social interaction time and sucrose preference for nonstressed B6 mice 2 weeks after the injections of PEI/*Gdnf* or PEI/*Egfp* complexes. We found that GDNF overexpression increased the social interaction time (Figure 1J), but not the sucrose preference (Figure 1K). We next investigated the effect of GDNF overexpression in stressful conditions. BALB mice were subjected to 4 weeks of CUMS and injected bilaterally into the NAc with either PEI/*Gdnf* or PEI/*Egfp* complexes on day 14 of the CUMS session. After the CUMS session, we performed behavioral assays. We found that the social interaction time (Figure 1J) and sucrose preference (Figure 1K) of the stressed BALB mice that received PEI/*Gdnf* complexes were significantly greater than those of the mice receiving PEI/*Egfp* complexes. These results suggest a crucial role for GDNF in social interactions and sucrose preference. The transcriptional regulation of *Gdnf* in the NAc may also be involved in the development of susceptibility and adaptation to CUMS.

Regulation of Histone Modifications by CUMS and Continuous IMI Treatment

To explore the molecular mechanisms by which CUMS alters *Gdnf* mRNA levels, resequence analysis of the *Gdnf* promoter (4000 base pairs) was performed on BALB and B6 mice. No differences were observed between the two mice strains (data not shown), suggesting that epigenetic regulations may account for altered *Gdnf* expression in stressed mice. Next, we measured the levels of several posttranslational histone modifications to the *Gdnf* promoter in vSTR tissues using a ChIP assay. We found several differences in the histone modifications of both BALB and B6 mice after CUMS and/or continuous IMI treatment. Q-PCR measurements indicated that *Gdnf* promoter-containing DNA fragments were significantly less common in the acetylated histone 3 (H3ac) immunoprecipitates prepared from stressed BALB mice. This effect was reversed by continuous IMI treatment (Figure 2A). Acetylated histone 4 (H4ac) levels at the *Gdnf* promoter were not affected by either CUMS or continuous IMI treatment (Figure 2B). In stressed B6 mice, H3ac levels at the *Gdnf* promoter, but not H4ac levels, were significantly increased by CUMS (Figures 2A and 2B). We also examined the effects of CUMS on the level of trimethylated histone 3 at lysine 27 (H3K27me3) and trimethylated histone 3 at lysine

4 (H3K4me3), which are the respective repressive and activating markers of transcription, at the *Gdnf* promoter. The levels of H3K27me3 were not affected by CUMS and IMI in BALB mice, but they were significantly reduced in B6 mice by CUMS (Figure 2C). The levels of H3K4me3 were significantly reduced by CUMS in both strains, and this reduction was reversed by IMI in stressed BALB mice (Figure 2D). These data suggest that histone modifications to the *Gdnf* promoter in response to CUMS are differentially regulated in each mouse strain.

Next, we investigated the mechanisms underlying the changes in the histone acetylation of the *Gdnf* promoter. We hypothesized that the altered expression of histone deacetylases (HDACs) could account for the altered level of histone acetylation. The levels of mRNA for HDACs (HDAC 1–11) were measured in the vSTR of BALB mice using Q-PCR. Several significant changes in *Hdacs* expression were observed following CUMS and/or continuous IMI treatment (Figure 2E). Of particular note, the mRNA level of *Hdac2* in stressed mice increased approximately two-fold compared with that of nonstressed controls. This enhancement was reversed by continuous IMI treatment. Changes at the protein level were also determined using Western blot analysis (Figure 2F). However, in the HP of BALB mice (Figure 2G) and the vSTR of B6 mice (Figure 2H), there were no significant effects of CUMS or IMI treatment on HDAC2 expression. Thus, these results suggest that HDAC2 may be an important regulator of the epigenetic repression of *Gdnf* expression in the vSTR of stressed BALB mice.

To determine whether CUMS influences the binding of HDAC2 to the *Gdnf* promoter, we performed a ChIP assay with vSTR DNA. Q-PCR measurements indicated that *Gdnf* promoter-containing DNA fragments are enriched in HDAC2 immunoprecipitates prepared from stressed BALB mice, and this effect was reversed by continuous IMI treatment (Figure 2I). No changes were observed at the *Bdnf* promoter II region (Figure S6A), whose transcript (*Bdnf* exon II) was not altered by either CUMS or IMI treatment (Figure S6B). This finding validates the specificity of the ChIP assay used in this study. In contrast to BALB mice, there was no significant effect of CUMS on HDAC2 binding to the *Gdnf* promoter in B6 mice (Figure 2J).

Rapid Antidepressant Effects of SAHA on CUMS-Induced Behavioral Deficits

Our data indicate that CUMS increases HDAC2 expression in the vSTR of BALB mice but not in B6 mice. This observation led to the hypothesis that this effect may be important for the transcriptional repression of *Gdnf* and the behavioral susceptibility to CUMS. To test the functional role of altered H3ac levels at the *Gdnf* promoter and HDAC2 expression in stressed BALB mice, suberoylanilide hydroxamic acid (SAHA), an HDAC inhibitor, was systemically administered (25 mg/kg/day) for the last 5 days of each 6-week CUMS sessions and during behavioral testing. In addition, to evaluate the possible antidepressant effects of SAHA, either IMI or fluoxetine (FLX), a selective serotonin reuptake inhibitor, was administered (25 mg/kg/day). The experimental design is shown in Figure S1C. The mice that received subchronic SAHA but not subchronic IMI or FLX




Figure 2. Differential Regulations of Histone Modifications in Stress-Susceptible and Stress-Adaptive Mice Strains

(A–D) Levels of posttranslational histone modifications in the *Gdnf* promoter of the vSTR of BALB and B6 mice subjected to CUMS or nonstressed (NS) conditions with or without IMI treatment were measured using ChIP assays with antibodies for acetylated histone 3 (A), acetylated histone 4 (B), and histone 3 trimethylated on lysine 27 (H3-K27 trimethylation) (C) or lysine 4 (H3-K4 trimethylation) (D) ($n = 6$ per group; * $p < 0.05$ versus NS mice receiving vehicle (normal water) in corresponding strains).

(E) mRNA levels of *Hdacs* in the vSTR of BALB mice subjected to CUMS or NS conditions with or without IMI treatment ($n = 6$ per group; * $p < 0.05$ versus NS mice receiving water).

(F and G) Western blot analysis of HDAC2 protein levels in the vSTR (F) and HP (G) of BALB mice subjected to CUMS or NS conditions with or without IMI treatment ($n = 7–8$ per group; * $p < 0.05$ versus NS mice receiving water).

(H) mRNA and protein levels of HDAC2 in the vSTR of stressed B6 mice ($n = 6$ per group).

(I and J) HDAC2 levels at the *Gdnf* promoter in the vSTR of BALB (I) and B6 (J) mice were measured using ChIP assays with a specific antibody for HDAC2 ($n = 7–8$ per group; * $p < 0.05$ versus NS mice receiving water). Data are presented as mean \pm SEM.

exhibited increased social interaction times compared with vehicle-treated mice in stressed conditions (Figure 3A). Similarly, the sucrose preference of mice receiving SAHA, but not IMI or FLX, was significantly increased compared to that of mice receiving vehicle in stressed conditions (Figure 3B). In the novelty-suppressed feeding test, SAHA reduced the latency to feed in mice from both the nonstressed and the stressed conditions, whereas subchronic IMI and FLX treatments did not affect the latency to feed (Figure 3C). In addition, the immobility times during the forced swim test were significantly decreased for mice receiving SAHA, but not IMI or FLX, compared to vehicle-treated mice from both the nonstressed and the stressed conditions (Figure 3D). Furthermore, subchronic SAHA treatment, but not IMI or FLX treatments, increased the mRNA levels of *Gdnf* in the vSTR of stressed mice (Figure 3E). These data suggest that HDAC inhibition can reverse both the increased depression-like behaviors and the reduction of *Gdnf* expression by CUMS. Our results also imply that SAHA has a more rapid antidepressant effect than IMI and FLX.

Role of HDAC2 in Behavioral Responses to CUMS

To test the direct contribution of HDAC2 in the NAc to CUMS-induced depression-like behaviors, dominant-negative HDAC2 (dnHDAC2; HDAC2 H141A) was overexpressed in the NAc of BALB mice using adeno-associated virus (AAV)-mediated gene transfer. Replacing His141 with Ala in the catalytic domain of HDAC2 reduces deacetylase activity by 75% (Humphrey et al., 2008). The experimental design is shown in Figure S1D. The successful transduction of AAV-mediated dnHDAC2 and control EGFP was first confirmed: EGFP fluorescence was observed in the NAc (Figure 3F), and Western blot analysis showed that dnHDAC2 was overexpressed in the vSTR region (Figure 3G). The NAc was then bilaterally infected with AAV-dnHDAC2 or AAV-EGFP. Seven days after the injection of AAV, mice were subjected to CUMS for 4 weeks, followed by the social interaction and sucrose preference tests. Mice that received AAV-dnHDAC2 exhibited increased social interaction times (Figure 3H) and sucrose preferences (Figure 3I) compared with the mice that received AAV-EGFP. Furthermore, the mRNA levels




Figure 3. Inhibition of HDAC2 Function Leads to a Stress-Resilient Phenotype

(A–E) Either SAHA, IMI, FLX, or saline were intraperitoneally administered (25 mg/kg of body weight for all drugs) on the last 5 days of each 6-week period of CUMS or nonstressed (NS) conditions and during behavioral testing. The social interaction times (n = 15–19 per group), (B) sucrose preferences (n = 18–20 per group), (C) latencies to feed (n = 18–20 per group), (D) immobility times (n = 18–20 per group), and (E) mRNA expression levels of *Gdnf* in the vSTR (n = 8 per group) are shown (*p < 0.05).

(F and G) Successful transductions of EGFP (F) and dominant-negative HDAC2 (dnHDAC2; G) using AAV-mediated gene transfer are shown.

(H–J) Effects of dnHDAC2 overexpression induced by AAV-mediated gene transfer in the NAc of BALB mice subjected to CUMS on (H) the social interaction time test (n = 14–15 per group), (I) the sucrose preference test (n = 17–19 per group), and (J) the *Gdnf* mRNA levels (n = 8 per group; *p < 0.05).

(K and L) Effects of the overexpression of wild-type HDAC2 (wtHDAC2) or the HDAC2 C262/274A mutant induced by AAV-mediated gene transfer in the NAc of B6 mice on the (K) social interaction time (n = 14–15 per group) and (L) mRNA levels of *Gdnf* (n = 8 per group; *p < 0.05). Data are presented as mean ± SEM.

of *Gdnf* in the vSTR of stressed mice that received AAV-dnHDAC2 were significantly increased compared to those of stressed mice injected with AAV-EGFP (Figure 3J). These results strongly suggest that the CUMS-induced activation of HDAC2 represses *Gdnf* transcription in the NAc, which results in aberrant behavioral responses in BALB mice.

To investigate the influence of HDAC2 on adaptive responses to CUMS in B6 mice, we overexpressed wild-type HDAC2 in the NAc of B6 mice and examined social interaction time and *Gdnf* expression. Stressed mice injected with AAV-HDAC2 did not show a reduction in social interaction time (Figure 3K) or *Gdnf* expression (Figure 3L) when compared with stressed mice injected with AAV-EGFP. A recent report showed that the nitrosylation of HDAC2 induces its release from chromatin, which promotes transcription. In the HDAC2 C262/274A mutant, which lacks S-nitrosylation sites, HDAC2 strongly associates with chromatin, thus repressing transcription (Nott et al., 2008). We investigated the effects of HDAC2 C262/274A overexpression

in the NAc of stressed B6 mice on social interaction and *Gdnf* expression. We found that stressed mice injected with AAV-HDAC2 C262/274A showed a reduction in social interaction time (Figure 3K) and *Gdnf* expression (Figure 3L) compared with stressed mice injected with AAV-EGFP. These results indicate that the gain of function of HDAC2 in B6 mice leads to a lack of active response to CUMS.

In contrast, the overexpression of the HDAC2 C262/274A mutant in nonstressed B6 mice did not affect the social interaction time or *Gdnf* expression (Figures 3K and 3L). Similar effects were also observed in nonstressed BALB mice receiving bilateral injections of either AAV-HDAC2 or AAV-HDAC2 C262/274A into the NAc (Figure S7). These manipulations did not alter the social interaction time (Figure S7B), sucrose preference (Figure S7C), or *Gdnf* expression (Figure S7D). These data suggest that other molecular mechanisms modulated by CUMS may also be involved in the HDAC2-mediated *Gdnf* repression and subsequent behavioral alterations.




Figure 4. CUMS Induces Hypermethylation of the *Gdnf* Promoter and Increases MeCP2 Binding to Its Promoter in Both Mice Strains

(A) Position of the CpG sites within the mouse *Gdnf* promoter.

(B) Methylation of the *Gdnf* promoter showing the frequency of methylation observed at each CpG site for the HP and vSTR (n = 8 per group; *p < 0.05).

(C) Mean percentages of the methylated clones for the HP and vSTR. The methylation percentage was calculated as the number of clones with at least one methylated CpG site divided by the total number of clones (n = 8 per group; *p < 0.05).

(D) mRNA levels of *Gdnf* in the HP, vSTR, dSTR, and prefrontal cortex (PFC) are shown (n = 6).

(E) CpG methylation profiles in the vSTR of stressed (CUMS) BALB mice with or without continuous IMI treatment, and nonstressed (NS) mice (n = 7–8 per group; *p < 0.05 versus NS mice receiving vehicle [normal water]).

(F) Samples of the sequence fluorograms obtained using bisulfite sequencing of DNA isolated from the vSTR of CUMS- and NS-BALB mice are shown. Arrows indicate methylated and nonmethylated sequences of CpG site 2.

(G) Mean percent of the methylation of CpG site 2 at the *Gdnf* promoter in stressed B6 mice (n = 7–8 per group; *p < 0.05).

(H) MeCP2 occupancy at the *Gdnf* promoter in the HP and vSTR were measured using ChIP analysis with antibodies specific to MeCP2 (n = 6 per group; *p < 0.05).

(I) MeCP2 levels at the *Gdnf* promoter and *Bdnf* promoter II were measured by ChIP analysis of vSTR DNA from mice subjected to CUMS (n = 7–8 per group; *p < 0.05 versus NS mice receiving water in the corresponding strain). Data are presented as mean \pm SEM.

CUMS Increases DNA Methylation at the *Gdnf* Promoter in Both Strains

Previous reports have suggested that histone methylation can affect DNA methylation at specific promoter regions (Lachner and Jenuwein, 2002). To investigate whether CUMS and/or IMI-induced alterations in the levels of H3K27me3 and H3K4me3 at the *Gdnf* promoter (Figures 2C and 2D) correlate with an increase in DNA methylation, DNA methylation assays

were performed. Cytosine methylation is a highly stable epigenetic process that regulates gene expression through its effects on transcription factor binding (Bird, 2001). Computational analysis (Takai and Jones, 2003) predicted that the *Gdnf* promoter has CpG islands adjacent to the transcription start site (CG > 60%, observed CpG/expected CpG > 0.81, and length > 300 bp; Figure 4A). Furthermore, these CpG islands are highly conserved in mice, rats, and humans (data not shown). First, to

examine whether CpG sites within the *Gdnf* promoter are truly methylated in vivo, the methylation levels of each CpG site were measured within the *Gdnf* promoter and a portion of the first exon. We used sodium bisulfite mapping to examine the methylation status of individual CpG sites within *Gdnf*. This method can detect both 5-methylcytosine and 5-hydroxy-methylcytosine. Sequence analysis of the bisulfite-converted DNA isolated from the HP and vSTR of BALB mice revealed less methylation at CpG sites 2 and 8–12 in the vSTR compared with congruent CpG sites in the HP (Figure 4B). In addition, sodium bisulfite mapping revealed a significantly lower percentage of methylated clones in the vSTR compared with the HP (Figure 4C). Concomitantly, the mRNA level of *Gdnf* in the vSTR was approximately 13-fold higher than that of the HP (Figure 4D), suggesting an association between the CpG methylation level and *Gdnf* mRNA expression in vivo. Therefore, the effects of 6 weeks of CUMS and continuous IMI treatment on CpG methylation were analyzed with bisulfite-converted DNA isolated from the vSTR of BALB mice. As indicated in Figures 4E and 4F, CUMS significantly increased methylation levels at CpG sites 2 and 3, but these hypermethylations were reversed by IMI treatment. Unexpectedly, the level of methylation at CpG site 2, but not at site 3, was also increased by CUMS in the vSTR of B6 mice (Figure 4G and data not shown).

CUMS Increases the Binding of MeCP2 at the *Gdnf* Promoter in Both Strains

The binding of methyl-CpG binding proteins (MBDs; MBD1, MBD2, MBD3, MBD4, and MeCP2) to the target gene promoter is a precise mechanism of gene transcription. Among MBDs, MeCP2 is most abundantly expressed as a chromosomal protein and requires a single methylated CpG site for preferential binding to DNA (Nan et al., 1997; Jones et al., 1998). Therefore, the binding of MeCP2 to the *Gdnf* promoter was directly assessed using the ChIP assay. First, to determine whether there is a difference in binding of MeCP2 to this promoter in the HP and vSTR of naive adult BALB mice, Q-PCR analysis of recovered DNA was performed using *Gdnf* promoter-specific primers. *Gdnf* promoter-containing DNA fragments were significantly less common in MeCP2 immunoprecipitates prepared from the vSTR compared with those from the HP (Figure 4H). Q-PCR analysis of the same immunoprecipitates was performed with a specific primer for *Gdnf* exon 3, which has no CpG island, and the immunoprecipitated DNA fragments were less common or undetectable (data not shown), validating the specificity of the ChIP protocol used. Next, the effect of 6 weeks of CUMS and continuous IMI treatment on the binding of MeCP2 to the *Gdnf* promoter was analyzed in the vSTR (Figure 4I). ChIP analysis revealed that CUMS significantly increased MeCP2 binding to the *Gdnf* promoter in both BALB and B6 mice, and continuous IMI treatment reversed this effect in stressed BALB mice. There was no significant difference in the binding of MeCP2 to the *Bdnf* promoter II region, which was assessed as a control. These results indicate that CUMS enhances the binding of MeCP2 to the *Gdnf* promoter in both mouse strains.

We next investigated the functional role of methylated CpG site 2 on *Gdnf* expression in Neuro2a cells. Treatment of these cells with 5-aza-2'-deoxycytidine, an inhibitor of DNA methylation, reduced the methylation level at the *Gdnf* promoter

(Figure S8A) and concomitantly increased *Gdnf* mRNA expression (Figure S8B). Next, the promoter activity of a CpG site 2-specific methylated *Gdnf* luciferase reporter gene was investigated. We found that CpG site 2-specific methylation resulted in an approximately 68% decrease in reporter activity when MeCP2 and HDAC2 were cotransfected into Neuro2a cells (Figure S8C). Previous reports have indicated that the high-affinity binding of MeCP2 to methylated DNA requires a run of four or more A/T bases adjacent to the methylated CpG site (Klose et al., 2005). We found two runs of A/T motifs located downstream of CpG site 2 (Figure S8D). To test the role of these motifs on *Gdnf* promoter activity, wild-type and mutant reporters were constructed for the A/T motifs in CpG site 2 (m1, m2, and m3; Figure S8D). Then, the promoter activity of the CpG site 2-specific methylated and nonmethylated luciferase reporters was measured using cotransfection experiments with MeCP2 and HDAC2 in Neuro2a cells (Figure S8E). We found that in nonmethylated conditions, there was no mutation effect on reporter activity by cotransfection with MeCP2 and HDAC2, whereas in the specific methylation of CpG site 2, the reporter activities of wild-type and m1 and m2 mutants, but not m3 mutant, were affected by HDAC2 and MeCP2 overexpression. These results suggest that the A/T motifs adjacent to CpG site 2 are critically involved in the MeCP2-HDAC2-mediated silencing of *Gdnf* transcription. Furthermore, we found that among the MBDs, MeCP2 was the most potent repressor of the CpG site 2-specific methylated reporter vector (Figure S8F). Together with the results observed in vivo, these findings suggest that the methylation of CpG site 2 is important for the epigenetic repression of *Gdnf* expression.

CUMS Increases the Binding of MeCP2-HDAC2 to the *Gdnf* Promoter in BALB Mice

The decreased expression level of *Gdnf* after CUMS in BALB mice was investigated to determine if it is triggered by the binding of MeCP2-HDAC2 complexes to the methylated CpG site of the *Gdnf* promoter. This hypothesis was supported, in part, by the finding that MeCP2 and HDAC2 are colocalized in the NAc (Figure 5A). The interactions of MeCP2 and HDAC2 were assessed using IP-Western blot analysis of vSTR proteins. We found that CUMS increased the formation of MeCP2-HDAC2 complexes in stressed BALB mice. This effect was reversed by continuous IMI treatment (Figure 5B). Next, to investigate the effect of CUMS on the binding of MeCP2-HDAC2 complexes at the *Gdnf* promoter, we performed re-ChIP assays using an antibody for HDAC2 on the vSTR samples that were initially immunoprecipitated with an antibody for MeCP2. The re-ChIP assays indicated that the *Gdnf* promoter-containing DNA fragments of stressed BALB mice, but not B6 mice, were significantly enriched compared with those of nonstressed mice, and this effect was reversed by continuous IMI treatment (Figure 5C). These results suggest that the CUMS-induced binding of MeCP2-HDAC2 complexes to the *Gdnf* promoter silences its transcription.

To investigate the role of DNA methylation in the CUMS-induced suppression of *Gdnf* expression and on depression-like behaviors, zebularine (ZEB), a DNA methyltransferase inhibitor, was continuously delivered into the NAc of BALB mice by an osmotic pump. The experimental design is shown in




Figure 5. Increased MeCP2-HDAC2 Occupancy at the *Gdnf* Promoter in Stressed BALB Mice

(A) Immunohistochemistry for HDAC2 and MeCP2 demonstrated strong nuclear staining in the NAc. Scale bar, 100 μ m.

(B) Nuclear extracts prepared from the vSTR of stressed (CUMS) or nonstressed (NS) BALB mice with or without IMI treatment were immunoprecipitated to evaluate the association of HDAC2 with MeCP2.

(C) Q-PCR assays of the reimmunoprecipitates of HDAC2 antibodies of ChIP samples treated with MeCP2 antibodies (re-ChIP assay) showed that *Gdnf* promoter-containing DNA fragments were enriched in stressed BALB mice ($n = 6-8$ per group; * $p < 0.05$ versus NS mice receiving vehicle). Data are presented as mean \pm SEM.

Figure S1E. Five days after surgery, mice were subjected to 4 weeks of CUMS, followed by behavioral and expression analyses. We found that the social interaction times and sucrose preferences of stressed mice receiving ZEB (100 μ M) were significantly higher compared with those times and preferences of vehicle-treated mice (Figures 6A and B). In the novelty-suppressed feeding test, the latency to feed was significantly decreased in stressed mice receiving ZEB compared with vehicle-treated controls (Figure 6C). In the forced swim test, the immobility times were significantly shorter in stressed and nonstressed mice receiving ZEB compared with the times of vehicle-treated mice (Figure 6D). Furthermore, the mRNA levels of *Gdnf* in ZEB-treated mice were greater than the levels in vehicle-treated mice (Figure 6E) in stressed conditions. These findings confirm that there is less DNA methylation of CpG site 2 at the *Gdnf* promoter in stressed mice treated with ZEB compared with vehicle-treated mice (Figure 6F). We also tested whether intra-NAc delivery of RG108, a potent, nonnucleoside inhibitor of DNA methylation, could reverse the increased depression-like behaviors in BALB mice. Similar to the effects of ZEB, continuous delivery of RG108 (100 μ M) directly into the NAc increased the social interaction time (Figure 6G) and sucrose preference (Figure 6H) of mice in the stressed condition. Furthermore, we found that CUMS increased the mRNA expressions for DNA methyltransferase 1 (DNMT1) and DNMT3a, but not DNMT3b, in the vSTR of stressed mice. This effect was reversed by continuous intra-NAc delivery of ZEB and RG108 (Figure 6I). These results suggest that DNA methylation is critical for the CUMS-induced *Gdnf* repression and subsequent depression-like behaviors in BALB mice. Our data also suggest that the continuous intra-NAc delivery of DNMT inhibitors represses the expression of Dnmts at the transcription level in postmitotic neurons.

CUMS Increases Binding of MeCP2-CREB to the *Gdnf* Promoter in B6 Mice

Although DNA methylation is generally thought to be associated with transcriptional repression of the target genes, a recent study

suggested that the binding of a complex of MeCP2 and cyclic AMP response element (CRE)-binding protein (CREB) to the methylated CpG site can activate transcription (Chahrouh et al., 2008). Interestingly, the putative CRE

site is adjacent to CpG site 2 of the *Gdnf* gene (Figure 7A). In addition, we found that MeCP2 and CREB are colocalized in the NAc (Figure 7B). These facts led us to speculate that the binding of the MeCP2-CREB complex to the *Gdnf* promoter may be a causal mechanism of the increased *Gdnf* expression in stressed B6 mice. To test this possibility, we assessed the interactions of MeCP2 and CREB in vSTR proteins of B6 and BALB mice. IP-Western blot analysis showed that there is no apparent difference in the formation of MeCP2-CREB complexes between stressed and nonstressed mice in both strains (Figure 7C). Next, to investigate the binding of MeCP2-CREB complexes at the *Gdnf* promoter, we performed re-ChIP assays using an antibody for CREB on vSTR samples that had been initially immunoprecipitated with an antibody for MeCP2. Consistent with a previous report (Chahrouh et al., 2008), CREB-MeCP2 complexes on the *somatostatin* promoter were enriched, whereas they were reduced on the *myocyte enhancer factor 2c* promoter (data not shown), validating the specificity of the re-ChIP used. We found that the *Gdnf* promoter-containing DNA fragments of stressed B6 mice were significantly enriched in the reimmunoprecipitates of samples treated with CREB antibodies compared with those of nonstressed mice. This effect was not seen in stressed BALB mice (Figure 7D). These results suggest that the CUMS-induced binding of MeCP2-CREB complexes to the *Gdnf* promoter leads to the activation of its transcription.

DISCUSSION

This study used genetically distinct inbred mouse strains to describe one of the molecular mechanisms underlying susceptibility and adaptation responses to chronic stress. The proposed mechanisms underlying stress susceptibility and adaptation are described in Figure 7E. Our results suggest that CUMS increases DNA methylation at CpG site 2, and this is associated with increased MeCP2 binding. MeCP2 associated with CpG site 2 interacts with HDAC2, which in turn decreases the level




Figure 6. Effects of a DNA Methyltransferase Inhibitor on CUMS-Induced Depression-Like Behaviors and *Gdnf* mRNA Expression

Either ZEB (10 μ M or 100 μ M) or a vehicle control was continuously and bilaterally delivered into the NAc of BALB mice. After each 4-week CUMS session, the depression-like behaviors of mice were analyzed.

(A–D) Social interaction times (n = 9–12 per group), (B) sucrose preferences (n = 10–12 per group), (C) latencies to feed (n = 10–12 per group), and (D) immobility times (n = 10–12 per group) are shown (*p < 0.05).

(E) The mRNA levels of *Gdnf* in the vSTR were measured by Q-PCR (n = 6–8 per group; *p < 0.05).

(F) Mean percent methylation of CpG site 2 at the *Gdnf* promoter in stressed BALB mice receiving ZEB (100 μ M) or vehicle (n = 6 per group; *p < 0.05).

(G and H) RG108 (10 μ M or 100 μ M) or vehicle was continuously and bilaterally delivered into the NAc of stressed BALB mice. After each 4-week CUMS session, the (G) social interaction time (n = 9–12 per group) and (H) sucrose preference (n = 11–15 per group) of the subjects were analyzed (*p < 0.05).

(I) The mRNA levels of *Dnmt1*, *Dnmt3a*, and *Dnmt3b* in the vSTR of mice receiving ZEB or RG108 (100 μ M) were measured by Q-PCR (n = 6–8 per group; *p < 0.05 versus NS mice receiving vehicle). Data are presented as mean \pm SEM.

of H3 acetylation and concomitantly represses *Gdnf* transcription, leading to the formation of a more depression-susceptible phenotype in BALB mice. Continuous IMI treatment relieves MeCP2 occupancy and reverses HDAC2 levels, which leads to normal levels of H3 acetylation and subsequent *Gdnf* transcription, resulting in normal emotional behaviors. Although increased DNA methylation at CpG site 2 and increased MeCP2 occupancy were also observed after CUMS exposure in B6 mice, the acetylation levels of H3 and *Gdnf* expression were greater. Importantly, we found evidence for the binding of the MeCP2-CREB complex to the methylated CpG site on the *Gdnf* promoter in stressed B6 mice. This may be a causal mechanism for the induction of *Gdnf* expression in stressed B6 mice. Thus, our data provide evidence that differential epigenetic marks in the

NAc, along with environmental and genetic factors, may influence either the susceptibility or adaptation responses of an organism to chronic daily stressful events.

Role of GDNF in Stress Responses

NAc has been implicated in the development of depression-like behaviors and has an influence on the action of antidepressants (Charney and Manji, 2004; Krishnan and Nestler, 2008; Feder et al., 2009). The data presented here indicate that differential histone modifications at the *Gdnf* promoter between stressed BALB and B6 mice result in differential levels of *Gdnf* expression. Overexpression of GDNF in the NAc increased social interaction times and sucrose preference in the stressed and/or the non-stressed conditions. Conditional GDNF knockout mice showed




Figure 7. Increased MeCP2-CREB Occupancy at the *Gdnf* Promoter in Stressed B6 Mice

(A) Positions of the CpG site 2 and putative CREB-binding site within the *Gdnf* promoter.

(B) Immunohistochemistry for MeCP2 and CREB demonstrated strong nuclear staining in the NAc. Scale bar, 100 μ m.

(C) Nuclear extracts prepared from the vSTR of stressed (CUMS) or non-stressed (NS) B6 mice were immunoprecipitated to evaluate the association of MeCP2 with CREB.

(D) Q-PCR analyses of reimmunoprecipitates for CREB antibodies of ChIP samples treated with MeCP2 antibodies (re-ChIP assay) showed that *Gdnf* promoter-containing DNA fragments were enriched in stressed B6 mice ($n = 4-5$ per group; $*p < 0.05$ versus NS mice). Data are presented as mean \pm SEM.

(E) Proposed mechanisms detailing how the chromatin microenvironment at the *Gdnf* promoter regulates its expression after CUMS exposure with or without IMI treatment in BALB and B6 mice. This study proposes that dynamic epigenetic changes in the *Gdnf* promoter may serve either as a repressive or activating marker of transcription in the NAc, and these changes may serve as causal mechanisms of the different behavioral responses to stress in BALB and B6 mice (refer to the Discussion for more details).

reduced spontaneous activity in the open field test (Pascual et al., 2008). In addition, mice that are not susceptible to social defeat stress show increased *Gdnf* expression in the ventral tegmental area (VTA) (Krishnan et al., 2007). The VTA-NAc network of the mesolimbic dopamine system may be involved in susceptibility and resistance responses to chronic stress (Nestler and Carlezon, 2006; Krishnan et al., 2007). GDNF promotes the survival and maintenance of midbrain dopamine-containing neurons, and GDNF protects neurons in the dopamine system from various toxic stimuli (Lin et al., 1993; Bespalov and Saarma, 2007; Pascual et al., 2008). Thus, the data presented here support the hypothesis that the mesolimbic dopamine system is involved in the formation of susceptibility and resistance responses to chronic stress.

In our experiments, continuous IMI treatment rescued the reduced GDNF expression in the vSTR of stressed BALB mice, suggesting that GDNF is also involved in the behavioral responses to antidepressants. The rescue of GDNF expression in stressed BALB mice returned behavioral performances back to control levels. However, it is still unclear whether the IMI-mediated upregulation of GDNF expression is critically involved in the antidepressant responses. IMI treatment also enhanced the mRNA expressions for other neurotrophic factors, including BDNF and VEGF, in multiple brain regions of BALB mice, and these molecules are thought to be associated with the behavioral responses to antidepressants (Warner-Schmidt and Duman, 2007; Krishnan and Nestler, 2008). Thus, we cannot exclude the possibility that molecules other than GDNF are important for the behavioral effects of antidepressant in the animal models used in this study. Further experiments are needed to clarify the role of GDNF in the behavioral responses to antidepressants.

CUMS and Antidepressants Affect Histone Modifications in the *Gdnf* Promoter

Persistent depressive symptoms suggest the involvement of stable changes in gene expression in brain, which may reflect a degree of chromatin remodeling, such as histone acetylation (Krishnan and Nestler, 2008; Tsankova et al., 2007). Recent reports have suggested that modulations of histone acetylation by HDAC2 and HDAC5 are also involved in the actions of antidepressants (Tsankova et al., 2006; Covington et al., 2009). In addition, subchronic administration of SAHA directly into the NAc of mice reverses the reduced social interaction time caused by social defeat stress (Covington et al., 2009). Similarly, this study demonstrated that the increased depression-like behaviors caused by CUMS were reversed by the subchronic administration of SAHA and the overexpression of dnHDAC2. However, nonstressed mice that received subchronic SAHA treatment did not exhibit any observable effects in their social interaction times, sucrose preferences, or expression levels of *Gdnf* mRNA. Taken together, these findings suggest that the hyperactive HDACs are involved in the reduction of *Gdnf* expression and subsequent depression-like behaviors induced by CUMS. In addition, we found that the overexpression of the HDAC2 C262/274A mutant, but not wild-type HDAC2, in the NAc of stressed B6 mice decreased social interaction time and *Gdnf* expression, suggesting a possible contribution of the S-nitrosylation of HDAC2 to the stress responses. We also found

that CUMS reduced the levels of H3K4me3 at the *Gdnf* promoter in both BALB and B6 mice, whereas the levels of H3K27me3 at its promoter were decreased only in B6 mice. These findings seem to be inconsistent with regard to the levels of *Gdnf* expression. The reduced H3K4me3 level at the *Gdnf* promoter in the NAc may be a common mechanism for responses to CUMS, and the reduced H3K27me3 level may be one of the important mechanisms modulating the chromatin microenvironment that primes adaptation responses to CUMS.

DNA Methylation at the *Gdnf* Gene Promoter Is Required for Both Susceptible and Adaptive Responses to CUMS

In addition to histone acetylation, the data presented here suggest an important role for DNA methylation in *Gdnf* expression and the subsequent behavioral responses to chronic stress. The epigenetic molecular mechanisms of DNA methylation in the brain may play important roles in the regulation of synaptic plasticity, memory formation, and stress responses (Weaver et al., 2004; Levenson and Sweatt, 2005; Krishnan and Nestler, 2008; Feder et al., 2009). Our data indicate that CUMS enhances DNA methylation at particular CpG sites on the *Gdnf* promoter in BALB mice. Importantly, our work indicates that the CUMS-induced depression-like behaviors and reduced *Gdnf* expression were reversed by the intra-NAc delivery of DNA methyltransferase inhibitors, a result that has been replicated in a recent report (LaPlant et al., 2010). Unexpectedly, the increased DNA methylation and MeCP2 binding also occurred in stress-resilient B6 mice. In general, DNA methylation is primarily associated with the repression of gene transcription. However, a recent study indicated that MeCP2-CREB complexes have assumed the role of inducing target gene expression (Chahrour et al., 2008). In addition, *Gdnf* expression may be regulated by CREB (Cen et al., 2006). Together with these findings, this study suggests that the binding of different MeCP2 complexes (i.e., MeCP2-CREB and MeCP2-HDAC2) to the methylated CpG site on the *Gdnf* promoter may be a causal mechanism for the induction and repression of *Gdnf* expression in the NAc of B6 and BALB mice.

Conclusion

This study provides insights into the role that genetic factors, in combination with environmental factors, may play in the epigenetic regulation of *Gdnf*. Dynamic epigenetic regulations of the *Gdnf* promoter in the NAc play important roles in determining both the susceptibility and the adaptation responses to chronic stressful events. Elucidation of the mechanisms underlying the modulations of HDAC2 expression, histone modifications, and DNA methylation influenced by CUMS could lead to novel approaches for the treatment of depression.

EXPERIMENTAL PROCEDURES

Details can be found in the Supplemental Experimental Procedures.

Animals

Adult male C57BL/6J and BALB/c mice (Charles River Japan) were maintained on a 12 hr/12 hr light/dark cycle with mouse chow and water ad libitum. Four mice were housed in each cage. Eight- or nine-week-old mice were used at the start of experiments (i.e., CUMS, stereotaxic surgery). All experimental

procedures were performed according to the Guidelines for Animal Care and Use at Yamaguchi University Graduate School of Medicine.

CUMS Procedure

The CUMS procedure has been previously described in detail (Lanfumeu et al., 1999; Rangon et al., 2007) and was conducted here with minor modifications. This procedure was based solely on environmental and social stressors, which did not include food/water deprivation. A total of three stressors were used in this study. For the first stressor, two of the following five ultra-mild diurnal stressors were delivered randomly over a period of 1–2 hr with a 2 hr stress-free time period between the two stressors: a period of cage tilt (30°), confinement to a small cage (11 × 8 × 8 cm), paired housing, soiled cage (50 ml water per 1 l of sawdust bedding), and odor (10% acetic acid). The second stressor consisted of four ultra-mild nocturnal stressors, including one overnight period with difficult access to food, one overnight period with permanent light, one overnight period with a 30° cage tilt, and one overnight period in a soiled cage. For the third stressor, a reversed light/dark cycle was used from Friday evening to Monday morning. This procedure was scheduled over a 1-week period and repeated four or six times, but the reversed light/dark cycle was omitted during the weekend of the last week (either the fourth or sixth week) of the session. Nonstressed mice were handled everyday for weighing purposes.

Behavioral Procedures

Behavioral tests were performed during the light phase (9 a.m. to 2 p.m.) with minor modifications, as reported previously (Uchida et al., 2008; 2010). All behavioral tests were conducted by experimenters who were blind to the treatment condition of the animal. Details can be found in the Supplemental Experimental Procedures.

Drugs

IMI, FLX, and 5-aza-2'-deoxycytidine were purchased from Sigma. ZEB and RG108 were purchased from Calbiochem. SAHA was synthesized as described previously (Suzuki et al., 2009). Details can be found in the Supplemental Experimental Procedures.

PEI-Mediated Gene Delivery

PEI-mediated gene delivery was performed as previously reported (Uchida et al., 2010). Plasmid DNA/PEI complexes were prepared according to the manufacturer's instructions (in vivo-jet PEI; PolyPlus Transfection). Seven days after bilateral canulae implantation into the NAc (+ 1.5 mm AP, ± 1.0 mm ML, –4.0 mm DV), mice were subjected to a 4-week CUMS session. PEI/plasmid complexes (0.5 μl/hemisphere) were injected on day 14 of the CUMS session. Details can be found in the Supplemental Experimental Procedures.

AAV-Mediated Gene Transfer

AAV-mediated gene transfer was performed as previously reported (Uchida et al., 2010). The genomic titer of each virus was determined using Q-PCR. The titers of AAV-EGFP, AAV-HA-HDAC2, AAV-HA-dnHDAC2, and AAV-HA-HDAC2 C262/274A were measured as 5.6×10^{12} viral genomes (vg)/ml, 3.1×10^{12} vg/ml, 3.5×10^{12} vg/ml, and 2.1×10^{12} vg/ml, respectively. For virus injections, the AAV vector (0.5 μl) was injected bilaterally into the NAc (+ 1.5 mm AP, ± 1.0 mm ML, –4.5 mm DV) at a rate of 0.1 μl/min. Mice were allowed to recover for 1 week after surgery. Details can be found in the Supplemental Experimental Procedures.

Statistical Analysis

Analyses of the data were performed using an appropriate analysis of variance. Significant effects were followed up with Bonferroni's post hoc tests. Unpaired t tests were used for two-group comparisons. Pearson correlations were calculated to assess correlations between data. In all cases, p values were two-tailed, and the comparisons were considered statistically significant when $p < 0.05$. Data are presented as the mean ± SEM.

SUPPLEMENTAL INFORMATION

Supplemental Information includes eight figures, four tables, and Supplemental Experimental Procedures and can be found with this article online at doi:10.1016/j.neuron.2010.12.023.

ACKNOWLEDGMENTS

The authors thank Drs. Koh Shinoda, Akinori Ishikawa, Yoichi Mizukami, Koh-ichi Udoh, and Tomoaki Murata for technical advice. This study was supported in part by a Grant-in-Aid for Scientific Research from the Japanese Ministry of Education, Culture, Sports, Science and Technology (S.U. and Y.W.) and a grant for Research on Psychiatric and Neurological Diseases and Mental Health from the Japanese Ministry of Health, Labor and Welfare (S.U. and Y.W.).

Accepted: October 28, 2010

Published: January 26, 2011

REFERENCES

- Berton, O., McClung, C.A., Dileone, R.J., Krishnan, V., Renthal, W., Russo, S.J., Graham, D., Tsankova, N.M., Bolanos, C.A., Rios, M., et al. (2006). Essential role of BDNF in the mesolimbic dopamine pathway in social defeat stress. *Science* 311, 864–868.
- Bespalov, M.M., and Saarna, M. (2007). GDNF family receptor complexes are emerging drug targets. *Trends Pharmacol. Sci.* 28, 68–74.
- Bhansali, P., Dunning, J., Singer, S.E., David, L., and Schmauss, C. (2007). Early life stress alters adult serotonin 2C receptor pre-mRNA editing and expression of the alpha subunit of the heterotrimeric G-protein G_q. *J. Neurosci.* 27, 1467–1473.
- Bird, A. (2001). Molecular biology: Methylation talk between histones and DNA. *Science* 294, 2113–2115.
- Cen, X., Nitta, A., Ohya, S., Zhao, Y., Ozawa, N., Mouri, A., Ibi, D., Wang, L., Suzuki, M., Saito, K., et al. (2006). An analog of a dipeptide-like structure of FK506 increases glial cell line-derived neurotrophic factor expression through cAMP response element-binding protein activated by heat shock protein 90/Akt signaling pathway. *J. Neurosci.* 26, 3335–3344.
- Chahrour, M., Jung, S.Y., Shaw, C., Zhou, X., Wong, S.T., Qin, J., and Zoghbi, H.Y. (2008). MeCP2, a key contributor to neurological disease, activates and represses transcription. *Science* 320, 1224–1229.
- Charney, D.S. (2004). Psychobiological mechanisms of resilience and vulnerability: implications for successful adaptation to extreme stress. *Am. J. Psychiatry* 161, 195–216.
- Charney, D.S., and Manji, H.K. (2004). Life stress, genes, and depression: multiple pathways lead to increased risk and new opportunities for intervention. *Sci. STKE* 2004, re5.
- Covington, H.E., 3rd, Maze, I., LaPlant, Q.C., Vialou, V.F., Ohnishi, Y.N., Berton, O., Fass, D.M., Renthal, W., Rush, A.J., 3rd, Wu, E.Y., et al. (2009). Antidepressant actions of histone deacetylase inhibitors. *J. Neurosci.* 29, 11451–11460.
- Duman, R.S., and Monteggia, L.M. (2006). A neurotrophic model for stress-related mood disorders. *Biol. Psychiatry* 59, 1116–1127.
- Feder, A., Nestler, E.J., and Charney, D.S. (2009). Psychobiology and molecular genetics of resilience. *Nat. Rev. Neurosci.* 10, 446–457.
- File, S.E., and Seth, P. (2003). A review of 25 years of the social interaction test. *Eur. J. Pharmacol.* 463, 35–53.
- Flavell, S.W., and Greenberg, M.E. (2008). Signaling mechanisms linking neuronal activity to gene expression and plasticity of the nervous system. *Annu. Rev. Neurosci.* 31, 563–590.
- Francis, D.D., Szegda, K., Campbell, G., Martin, W.D., and Insel, T.R. (2003). Epigenetic sources of behavioral differences in mice. *Nat. Neurosci.* 6, 445–446.
- Fyffe, S.L., Neul, J.L., Samaco, R.C., Chao, H.T., Ben-Shachar, S., Moretti, P., McGill, B.E., Goulding, E.H., Sullivan, E., Tecott, L.H., and Zoghbi, H.Y. (2008). Deletion of *Mecp2* in *Sim1*-expressing neurons reveals a critical role for MeCP2 in feeding behavior, aggression, and the response to stress. *Neuron* 59, 947–958.
- Hovatta, I., Tennant, R.S., Helton, R., Marr, R.A., Singer, O., Redwine, J.M., Ellison, J.A., Schadt, E.E., Verma, I.M., Lockhart, D.J., and Barlow, C. (2005). Glyoxalase 1 and glutathione reductase 1 regulate anxiety in mice. *Nature* 438, 662–666.
- Humphrey, G.W., Wang, Y.H., Hirai, T., Padmanabhan, R., Panchision, D.M., Newell, L.F., McKay, R.D., and Howard, B.H. (2008). Complementary roles for histone deacetylases 1, 2, and 3 in differentiation of pluripotent stem cells. *Differentiation* 76, 348–356.
- Jakobsson, J., Cordero, M.I., Bisaz, R., Groner, A.C., Buskamp, V., Bensadoun, J.C., Cammas, F., Losson, R., Mansuy, I.M., Sandi, C., and Trono, D. (2008). KAP1-mediated epigenetic repression in the forebrain modulates behavioral vulnerability to stress. *Neuron* 60, 818–831.
- Jones, P.L., Veenstra, G.J., Wade, P.A., Vermaak, D., Kass, S.U., Landsberger, N., Strouboulis, J., and Wolffe, A.P. (1998). Methylated DNA and MeCP2 recruit histone deacetylase to repress transcription. *Nat. Genet.* 19, 187–191.
- Klose, R.J., Sarraf, S.A., Schmiedeberg, L., McDermott, S.M., Stancheva, I., and Bird, A.P. (2005). DNA binding selectivity of MeCP2 due to a requirement for A/T sequences adjacent to methyl-CpG. *Mol. Cell* 19, 667–678.
- Krishnan, V., Han, M.H., Graham, D.L., Berton, O., Renthal, W., Russo, S.J., LaPlant, Q., Graham, A., Lutter, M., Lagace, D.C., et al. (2007). Molecular adaptations underlying susceptibility and resistance to social defeat in brain reward regions. *Cell* 131, 391–404.
- Krishnan, V., and Nestler, E.J. (2008). The molecular neurobiology of depression. *Nature* 455, 894–902.
- Lachner, M., and Jenuwein, T. (2002). The many faces of histone lysine methylation. *Curr. Opin. Cell Biol.* 14, 286–298.
- Lanfume, L., Pardon, M.C., Laaris, N., Joubert, C., Hanoun, N., Hamon, M., and Cohen-Salmon, C. (1999). 5-HT_{1A} autoreceptor desensitization by chronic ultramild stress in mice. *Neuroreport* 10, 3369–3374.
- LaPlant, Q., Vialou, V., Covington, H.E., III, Dumitriu, D., Feng, J., Warren, B.L., Maze, I., Dietz, D.M., Watts, E.L., Iñiguez, S.D., et al. (2010). *Dnmt3a* regulates emotional behavior and spine plasticity in the nucleus accumbens. *Nat. Neurosci.* 13, 1137–1143.
- Leonardo, E.D., and Hen, R. (2008). Anxiety as a developmental disorder. *Neuropsychopharmacology* 33, 134–140.
- Levenson, J.M., and Sweatt, J.D. (2005). Epigenetic mechanisms in memory formation. *Nat. Rev. Neurosci.* 6, 108–118.
- Lin, L.F., Doherty, D.H., Lile, J.D., Bektesh, S., and Collins, F. (1993). GDNF: a glial cell line-derived neurotrophic factor for midbrain dopaminergic neurons. *Science* 260, 1130–1132.
- Mozhui, K., Karlsson, R.M., Kash, T.L., Ihne, J., Norcross, M., Patel, S., Farrell, M.R., Hill, E.E., Graybeal, C., Martin, K.P., et al. (2010). Strain differences in stress responsivity are associated with divergent amygdala gene expression and glutamate-mediated neuronal excitability. *J. Neurosci.* 30, 5357–5367.
- Nan, X., Campoy, F.J., and Bird, A. (1997). MeCP2 is a transcriptional repressor with abundant binding sites in genomic chromatin. *Cell* 88, 471–481.
- Nestler, E.J., Barrot, M., DiLeone, R.J., Eisch, A.J., Gold, S.J., and Monteggia, L.M. (2002). Neurobiology of depression. *Neuron* 34, 13–25.
- Nestler, E.J., and Carlezon, W.A., Jr. (2006). The mesolimbic dopamine reward circuit in depression. *Biol. Psychiatry* 59, 1151–1159.
- Nott, A., Watson, M., Robinson, J.D., Crepaldi, L., and Riccio, A. (2008). S-nitrosylation of histone deacetylase 2 induces chromatin remodeling in neurons. *Nature* 455, 411–415.
- Palumbo, M.L., Zorrilla Zubieta, M.A., Cremaschi, G.A., and Genaro, A.M. (2009). Different effect of chronic stress on learning and memory in BALB/c and C57BL/6 inbred mice: involvement of hippocampal NO production and PKC activity. *Stress* 12, 350–361.

- Pascual, A., Hidalgo-Figueroa, M., Piruat, J.I., Pintado, C.O., Gomez-Diaz, R., and Lopez-Barneo, J. (2008). Absolute requirement of GDNF for adult catecholaminergic neuron survival. *Nat. Neurosci.* 11, 755–761.
- Porsolt, R.D., Le Pichon, M., and Jalfre, M. (1977). Depression: a new animal model sensitive to antidepressant treatments. *Nature* 266, 730–732.
- Rangon, C.M., Fortes, S., Lelievre, V., Leroux, P., Plaisant, F., Joubert, C., Lanfumey, L., Cohen-Salmon, C., and Gressens, P. (2007). Chronic mild stress during gestation worsens neonatal brain lesions in mice. *J. Neurosci.* 27, 7532–7540.
- Richardson-Jones, J.W., Craige, C.P., Guiard, B.P., Stephen, A., Metzger, K.L., Kung, H.F., Gardier, A.M., Dranovsky, A., David, D.J., Beck, S.G., et al. (2010). 5-HT1A autoreceptor levels determine vulnerability to stress and response to antidepressants. *Neuron* 65, 40–52.
- Suzuki, N., Suzuki, T., Ota, Y., Nakano, T., Kurihara, M., Okuda, H., Yamori, T., Tsumoto, H., Nakagawa, H., and Miyata, N. (2009). Design, synthesis, and biological activity of boronic acid-based histone deacetylase inhibitors. *J. Med. Chem.* 52, 2909–2922.
- Takai, D., and Jones, P.A. (2003). The CpG island searcher: a new WWW resource. *In Silico Biol.* 3, 235–240.
- Tsankova, N., Renthal, W., Kumar, A., and Nestler, E.J. (2007). Epigenetic regulation in psychiatric disorders. *Nat. Rev. Neurosci.* 8, 355–367.
- Tsankova, N.M., Berton, O., Renthal, W., Kumar, A., Neve, R.L., and Nestler, E.J. (2006). Sustained hippocampal chromatin regulation in a mouse model of depression and antidepressant action. *Nat. Neurosci.* 9, 519–525.
- Uchida, S., Hara, K., Kobayashi, A., Funato, H., Hobara, T., Otsuki, K., Yamagata, H., McEwen, B.S., and Watanabe, Y. (2010). Early life stress enhances behavioral vulnerability to stress through the activation of REST4-mediated gene transcription in the medial prefrontal cortex of rodents. *J. Neurosci.* 30, 15007–15018.
- Uchida, S., Nishida, A., Hara, K., Kamemoto, T., Suetsugi, M., Fujimoto, M., Watanuki, T., Wakabayashi, Y., Otsuki, K., McEwen, B.S., and Watanabe, Y. (2008). Characterization of the vulnerability to repeated stress in Fischer 344 rats: possible involvement of microRNA-mediated down-regulation of the glucocorticoid receptor. *Eur. J. Neurosci.* 27, 2250–2261.
- Warner-Schmidt, J.L., and Duman, R.S. (2007). VEGF is an essential mediator of the neurogenic and behavioral actions of antidepressants. *Proc. Natl. Acad. Sci. USA* 104, 4647–4652.
- Wong, M.L., and Licinio, L. (2001). Research and treatment approaches to depression. *Nat. Rev. Neurosci.* 2, 343–351.
- Weaver, I.C., Cervoni, N., Champagne, F.A., D'Alessio, A.C., Sharma, S., Seckl, J.R., Dymov, S., Szyf, M., and Meaney, M.J. (2004). Epigenetic programming by maternal behavior. *Nat. Neurosci.* 7, 847–854.